abcam

ab314228

Anti-CD204 antibody [EPR28270-88] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR28270-88
Purification technique	Affinity purification Protein A
Concentration	1.01 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse
Dilution info	-

Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	-
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	-
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	-
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

Target data

See full target information Msr1

Function	Membrane glycoproteins implicated in the pathologic deposition of cholesterol in arterial walls during atherogenesis. Two types of receptor subunits exist. These receptors mediate the endocytosis of a diverse group of macromolecules, including modified low density lipoproteins (LDL).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD204 also known as MSR1 or the macrophage scavenger receptor 1 is a class A scavenger receptor. It has a molecular mass of approximately 80-85 kDa. CD204 is mainly expressed on the surface of macrophages as well as some dendritic cells. This receptor plays a role in mediating the endocytosis of various polyanionic molecules such as modified low-density lipoproteins. CD204 serves as a recognition site for apoptotic cells signaling molecules and some types of bacteria.
Biological function summary	CD204 influences inflammatory responses and lipid metabolism. It does not function alone but as part of a complex involving several other molecules like pattern recognition receptors. CD204 recognizes and binds to modified lipoproteins and is significant in processes like phagocytosis and clearance of cellular debris. Its activity affects immune response modulation and can influence atherosclerosis development by its interaction with oxidized low-density lipoproteins.
Pathways	The role of CD204 extends to key cellular and immunological pathways. This receptor is involved in the Toll-like receptor signaling pathway which is important for innate immunity besides participating in the macrophage activation pathways. In these pathways CD204 interacts with other proteins including CD36 and SRA.

This interaction helps coordinate inflammatory responses and the resolution of inflammation.

Associated diseases and disorders

CD204 is associated with atherosclerosis and chronic inflammation. Its connection with these conditions arises from its role in lipid uptake and immune cell regulation within the vascular system. In the context of atherosclerosis CD204's interaction with Apolipoprotein E (ApoE) influences the formation of foam cells which are significant in the disease's progression. Similarly in chronic inflammation CD204 modulates macrophage behavior impacting the inflammatory environment and potentially affecting the disease outcome.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


10 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)


This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded Mouse lung cancer tissue labeling CD204 with ab314227 at 1/5000 (0.103 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on immune cells in mouse lung cancer.The

section was incubated with ab314227 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded Mouse neuroendocrine carcinoma tissue labeling CD204 with ab314227 at 1/5000 (0.103 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on immune cells in mouse neuroendocrine carcinoma.The section was incubated with ab314227 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded Mouse lung tissue labeling CD204 with ab314227 at 1/5000 (0.103 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on macrophages in mouse lung.The section was incubated with ab314227 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded (A) RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) cell pellets; (B) Neuro-2a (mouse neuroblastoma neuroblast) cell pellets. tissue labeling CD204 with ab314227 at 1/2000 (0.257 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on (A) RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) cell pellets, no staining on (B) Neuro-2a (mouse neuroblastoma neuroblast) cell pellets.The section was incubated with ab314227 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labeling CD204 with ab314227 at 1/5000 (0.103 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on immune cells in mouse colon.The section was incubated with ab314227 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse bone marrow cells labelling CD204 with ab314227 at 1/50 (10.26 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green). Confocal image showing positive staining in mouse bone marrow treated with lipopolysaccharide (2 ug/mL) for 24 h.Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). Anti-F4/80 rat monoclonal antibody was used to counterstain tubulin at 1/100 10 ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Immunocytochemistry/ Immunofluorescence - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)


This data was developed using ab314227, the same antibody clone in a different buffer formulation. Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) cells labelling CD204 with ab314227 at 1/50 (10.26 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green). Confocal image showing cytoplasmic staining in RAW 264.7 cell line.Negative control: Neuro-2a Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Flow Cytometry - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)


This data was developed using ab314227, the same antibody clone in a different buffer formulation. Flow cytometric analysis of Neuro-2a (mouse neuroblastoma neuroblast, Left) / Raw264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage, Right) cells labelling CD204 with ab314227 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with

primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Negative control: Neuro-2a.



Flow Cytometry - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Flow cytometric analysis of Isotype (Left) / Mouse bone marrow treated with 2ug/ml LPS for 24h (Middle) / Untreated mouse bone marrow (Right) cells labelling CD204 with ab314227 at 1/5000 dilution (0.01 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Cells are co-stained with F4/80 conjugated with Alexa Fluor®647. Gated on viable cells.



Western blot - Anti-CD204 antibody [EPR28270-88] - BSA and Azide free (ab314228)

This data was developed using ab314227, the same antibody clone in a different buffer formulation. Blocking and diluting buffer and concentration: 5% NFDM/TBST Negative control: Neuro-2a, NIH/3T3. The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 35937296). In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution. Exposure time: Lane 1: 5.5 seconds; Lane 2-4: 37 seconds.
All lanes:
Western blot - Anti-CD204 antibody [EPR28270-88] (ab314227) at 1/1000 dilution

Lane 1:
J774A.1 (mouse reticum cell sarcoma monocyte/macrophage) whole cell lysate at 20 µg

Lane 2:
RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate at 20 µg

Lane 3:
Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Lane 4:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 80 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.