abcam

ab314295

PE Anti-SIRP alpha antibody [EPR22930-163]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	PE
Excitation/Emission	Ex: 480;565nm, Em: 578nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR22930-163
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	1/2500

Notes	-

ICC/IF

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information SIRPA

Function

Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function (By similarity). Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Plays a role in antiviral immunity and limits new world arenavirus infection by decreasing virus internalization (By similarity). Receptor for THBS1 (PubMed:24511121). Interaction with THBS1 stimulates phosphorylation of SIRPA (By similarity). In response to THBS1, involved in ROS signaling in non-phagocytic cells, stimulating NADPH oxidase-derived ROS production (PubMed:24511121).

Storage

Shipped at conditions	Blue Ice

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	SIRP alpha also known as CD172a is a transmembrane receptor protein with a mass ranging between 70-110 kDa due to glycosylation. This protein extensively expresses on the surface of myeloid cells neurons and a subset of T-cells and is part of the immunoglobulin superfamily. SIRP alpha interacts with its ligand CD47 a widely expressed glycoprotein involved in immune response regulation. Its mechanical action primarily involves signal regulation through the recruitment of SHP-1 and SHP-2 two cytoplasmic tyrosine phosphatases.
Biological function summary	SIRP alpha functions significantly in the regulation of phagocytosis acting as a "don't eat me" signal to macrophages upon binding with CD47. It does not act alone; rather it is part of a complex that recruits SHP-1 and SHP-2 leading to inhibition of dephosphorylation activities essential for engulfment processes. This regulatory mechanism is important for maintaining cellular homeostasis ensuring that healthy cells are not mistakenly destroyed by the immune system.
Pathways	SIRP alpha plays an important role in the innate immune pathways involving the regulation of phagocytosis and cell-cell adhesion. Particularly it fits into the

	immune checkpoint pathways where it interacts closely with proteins like CD47 and plays a role in the interaction between the immune system and cancer cells. Through these pathways SIRP alpha helps maintain balance in the immune response allowing for the recognition of self versus non-self therefore preventing autoimmunity while facilitating the clearance of pathogens.
Associated diseases and disorders	SIRP alpha is implicated in cancer and autoimmune diseases. In cancer its interaction with CD47 allows tumor cells to escape phagocytosis promoting tumor survival and growth. In autoimmune disorders dysregulated SIRP alpha expression or signaling could miscommunicate immune signals leading to the destruction of healthy tissue. Understanding the link between SIRP alpha and these conditions can reveal potential targets for therapeutic development especially using inhibitors or modulators targeting the SIRP alpha-CD47 interaction.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - PE Anti-SIRP alpha antibody [EPR22930-163] (ab314295)

SIRP alpha Flow Cytometry staining using rabbit Anti-SIRP alpha antibody

Flow cytometry overlay histogram showing left THP-1 positive cells and right negative Jurkat stained with ab314295 (red line). The cells were incubated in 1x PBS containing 10µg/ml human IgG and 10% normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab314295) (1x 10^6 in 100µl at 0.2 µg/ml (1/2500)) for 30min on ice.

Isotype control antibody (black line) was Rabbit IgG (monoclonal) Phycoerythrin Isotype Control (ab209478) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5000 events were collected using a 50 mW Yellow/Green laser (561nm) and 585/42 bandpass filter.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.