abcam

ab314504

Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker

Anti-SQSTM1 / p62 antibody [RM1079] (ab314504) is a rabbit recombinant multiclonal antibody detecting SQSTM1 / p62 in **Western Blot, IP, IHC-P, IHC-Fr, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency

Lab Essentials | Recombinant | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Multiclonal
Immunogens	The exact immunogen used to generate this antibody is proprietary information.
Clone number	RM1079
Purification technique	Affinity purification Protein A
Concentration	0.524 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/1000 - 1/20000
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ICC/IF

Tested

Species	Human
Dilution info	1/500

Notes	-

Species	Mouse
Dilution info	1/500
Notes	-

Species	Rat
Dilution info	1/500
Notes	-

Flow Cyt (Intra)

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse

Dilution info	1/50
Notes	-

Species	Rat
Dilution info	1/50
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information SQSTM1

Function

Molecular adapter required for selective macroautophagy (aggrephagy) by acting as a bridge between polyubiquitinated proteins and autophagosomes (PubMed:15340068, PubMed:15953362, PubMed:16286508, PubMed:17580304, PubMed:20168092, PubMed:22017874, PubMed:22622177, PubMed:24128730, PubMed:28404643, PubMed:29343546, PubMed:29507397, PubMed:31857589, PubMed:33509017, PubMed:34471133, PubMed:34893540, PubMed:35831301, PubMed:37306101, PubMed:37802024). Promotes the recruitment of ubiquitinated cargo proteins to autophagosomes via multiple domains that bridge proteins and organelles in different steps (PubMed:16286508, PubMed:20168092, PubMed:22622177, PubMed:24128730, PubMed:28404643, PubMed:29343546, PubMed:29507397, PubMed:34893540, PubMed:37802024). SQSTM1 first mediates the assembly and removal of ubiquitinated proteins by undergoing liquid-liquid phase separation upon binding to ubiquitinated proteins via its UBA domain, leading to the formation of insoluble cytoplasmic inclusions, known as p62 bodies (PubMed:15911346, PubMed:20168092, PubMed:22017874, PubMed:24128730, PubMed:29343546, PubMed:29507397, PubMed:31857589, PubMed:37802024). SQSTM1 then interacts with ATG8 family proteins on autophagosomes via its LIR motif, leading to p62 body recruitment to autophagosomes, followed by autophagic clearance of ubiquitinated proteins (PubMed:16286508, PubMed:17580304, PubMed:20168092, PubMed:22622177, PubMed:24128730, PubMed:28404643, PubMed:37802024). SQSTM1 is itself degraded along with its ubiquitinated cargos (PubMed:16286508, PubMed:17580304, PubMed:37802024). Also required to recruit ubiquitinated proteins to PML bodies in the nucleus (PubMed:20168092). Also involved in autophagy of peroxisomes (pexophagy) in response to reactive oxygen species (ROS) by acting as a bridge between ubiquitinated PEX5 receptor and autophagosomes (PubMed:26344566). Acts as an activator of the NFE2L2/NRF2 pathway via interaction with KEAP1: interaction inactivates the BCR(KEAP1) complex by sequestering the complex in inclusion bodies, promoting nuclear accumulation of NFE2L2/NRF2 and subsequent

expression of cytoprotective genes (PubMed:20452972, PubMed:28380357, PubMed:33393215, PubMed:37306101). Promotes relocalization of 'Lys-63'-linked ubiquitinated STING1 to autophagosomes (PubMed:29496741). Involved in endosome organization by retaining vesicles in the perinuclear cloud: following ubiquitination by RNF26, attracts specific vesicle-associated adapters, forming a molecular bridge that restrains cognate vesicles in the perinuclear region and organizes the endosomal pathway for efficient cargo transport (PubMed:27368102, PubMed:33472082). Sequesters tensin TNS2 into cytoplasmic puncta, promoting TNS2 ubiquitination and proteasomal degradation (PubMed:25101860). May regulate the activation of NFKB1 by TNF, nerve growth factor (NGF) and interleukin-1 (PubMed:10356400, PubMed:10747026, PubMed:11244088, PubMed:12471037, PubMed:16079148, PubMed:19931284). May play a role in titin/TTN downstream signaling in muscle cells (PubMed:15802564). Adapter that mediates the interaction between TRAF6 and CYLD (By similarity).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-SQSTM1 / p62 antibody [RM1079] (ab314504) is a rabbit recombinant recombinant multiclonal antibody and is validated for use in Western Blot (WB), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunohistochemistry (IHC-Fr), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of SQSTM1 / p62?
Anti-SQSTM1 / p62 [RM1079] (ab314504) specifically detects a band for SQSTM1 / p62 (UniProt: Q13501) at a molecular weight of 47kDa.

Specificity confirmed
The specificity of Anti-SQSTM1 / p62 antibody [RM1079] (ab314504) has been confirmed by Western blot testing in SQSTM1 Knockout HAP1 cells.

What are recombinant multiclonals?
Recombinant multiclonals are a mixture of recombinant antibodies co-expressed from a library of heavy and light chains. They offer several advantages including:

- The sensitivity of polyclonal antibodies by recognising multiple epitopes
- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

View our range of recombinant multiclonal antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

13 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human breast cancer using rabbit Anti-SQSTM1 / p62 antibody

Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue labeling SQSTM1 with ab314504 at 1/5000 (0.101 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human breast cancer.The section was incubated with ab314504 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins


---



Immunocytochemistry/ Immunofluorescence - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SQSTM1 / p62 antibody


Immunofluorescent analysis of 100% methanol-fixed, 0.1% TritonX-100 permeabilized SQSTM1 KO HAP1(SQSTM1 knockout human chronic myelogenous leukemia cell line) cells labelling SQSTM1 with ab314504 at 1/500 (1.01 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2ug/ml dilution (Green). Confocal image showing cytoplasmic staining in HAP1 cell line, and negative staining in SQSTM1 KO HAP1 cell line. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2ug/ml dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SQSTM1 / p62 antibody

Immunohistochemical analysis of paraffin-embedded (A) Wild-type HCT116 (human colon epithelial) cell pellet (B) SQSTM1 knockout HCT116 (ab266871) cell pellet tissue labeling SQSTM1 with ab314504 at 1/20000 (0.025 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on (A) Wild-type HCT116 cell pellet, no staining on (B) SQSTM1 knockout HCT116 (ab266871) cell pellet. The section was incubated with ab314504 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins



Immunoprecipitation - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)


SQSTM1 was immunoprecipitated from 0.35 mg HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate with ab314504 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab314504 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution. Lane 1: HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate Lane 2: ab314504 IP in HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab314504 in HepG2 whole cell lysate Blocking and dilution buffer and concentration: 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504) at 1/30 dilution

All lanes:
HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution


Exposure time: 62s



Immunocytochemistry/ Immunofluorescence - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunocytochemistry/ Immunofluorescence staining of PC-12 (rat adrenal gland pheochromocytoma cell) using rabbit Anti-SQSTM1 / p62 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized PC-12 (rat adrenal gland pheochromocytoma cell) cells labelling SQSTM1 with ab314504 at 1/500 (1.01 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2ug/ml dilution (Green). Confocal image showing cytoplasmic staining in PC-12 cell line. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2ug/ml dilution.




Immunocytochemistry/ Immunofluorescence - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunocytochemistry/ Immunofluorescence staining of MEF (mouse embryo fibroblast cell) using rabbit Anti-SQSTM1 / p62 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized MEF (mouse embryo fibroblast cell) cells labelling SQSTM1 with ab314504 at 1/500 (1.01 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2ug/ml dilution (Green). Confocal image showing cytoplasmic staining in MEF cell line. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2ug/ml dilution.



Immunohistochemistry (Frozen sections) - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunohistochemistry (Frozen sections) staining of Rat cerebrum (fresh) using rabbit Anti-SQSTM1 / p62 antibody


Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat cerebrum (fresh) tissue labeling SQSTM1 with ab314504 at 1/50 (10.1 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Confocal image showing positive staining on rat cerebrum. The nuclear counterstain was DAPI (Blue). The section was incubated with ab314504 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse cerebrum using rabbit Anti-SQSTM1 / p62 antibody


Immunohistochemical analysis of paraffin-embedded Mouse cerebrum tissue labeling SQSTM1 with ab314504 at 1/1000 (0.505 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse cerebrum.The section was incubated with ab314504 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins



Immunohistochemistry (Frozen sections) - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunohistochemistry (Frozen sections) staining of Mouse cerebrum (fresh) using rabbit Anti-SQSTM1 / p62 antibody


Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse cerebrum (fresh) tissue labeling SQSTM1 with ab314504 at 1/50 (10.1 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Confocal image showing positive staining on mouse cerebrum. The nuclear counterstain was DAPI (Blue). The section was incubated with ab314504 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Rat cerebrum using rabbit Anti-SQSTM1 / p62 antibody

Immunohistochemical analysis of paraffin-embedded Rat cerebrum tissue labeling SQSTM1 with ab314504 at 1/1000 (0.505 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat cerebrum.The section was incubated with ab314504 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins



Western blot - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Western blot staining using rabbit Anti-SQSTM1 / p62 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST Exposure time: Lane 1: 103 seconds; Lane 2 and 3: 125 seconds
All lanes:
Western blot - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504) at 1/1000 dilution

Lane 1:
Human kidney tissue lysate at 20 µg


Lane 2:
Mouse spleen tissue lysate at 20 µg

Lane 3:
Rat spleen tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 62 kDa



Western blot - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Western blot staining using rabbit Anti-SQSTM1 / p62 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST Exposure time: Lanes 1-4: 15 seconds; Lanes 5-8: 103 seconds
All lanes:
Western blot - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504) at 1/1000 dilution

Lane 1:
MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
293T (human embryonic kidney epithelial cell) whole cell lysate at 20 µg

Lane 3:
K-562 (human chronic myelogenous leukemia lymphoblast) whole cell lysate at 20 µg

Lane 4:
MEF (mouse embryo fibroblast) whole cell lysate at 20 µg

Lane 5:
C6 (rat glial tumor glial cell) whole cell lysate at 20 µg

Lane 6:
RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate at 20 µg


Lane 7:
PC-12 (rat adrenal gland pheochromocytoma cell) whole cell lysate at 20 µg

Lane 8:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 62 kDa



Western blot - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504)

SQSTM1 / p62 Western blot staining using rabbit Anti-SQSTM1 / p62 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST The samples were run on a Bis-Tris gel under reducing conditions. Western blot: Anti-SQSTM1 / p62 antibody (ab314504) staining at 1/1000 dilution, shown in green; Mouse Anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20, 000 dilution, shown in red. In Western blot, ab314504 was shown to bind specifically to SQSTM1 / p62. Target of interest was observed at 62 kDa in wild-type HAP1 cell lysates (lane 1) with no signal observed at this size in SQSTM1 knockout cell line (lane 2). To generate this image, samples were first run on an SDS-PAGE gel then transferred onto an immobilon-FL PVDF membrane. Membranes were blocked in a fluorescent western blot (TBS-based) blocking solution before incubation with primary antibodies overnight at 4 °C. Blots were washed in TBS-T, incubated with secondary antibodies Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution for 1 h at room temperature, washed again then imaged.
All lanes:
Western blot - Anti-SQSTM1 / p62 antibody [RM1079] - Autophagosome Marker (ab314504) at 1/1000 dilution

Lane 1:
Wild type HAP1 (human chronic myelogenous leukemia cell) whole cell lysate at 20 µg

Lane 2:
SQSTM1 knockout HAP1 whole cell lysate at 20 µg

Lane 3:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 4:

HepG2 (human hepatocellar carcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG H&L (800CW) and Goat Anti-Mouse IgG H&L (680RD) at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 62 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.