abcam

ab314864

Anti-HMGN1 antibody [EPR26772-17]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR26772-17
Purification technique	Affinity purification Protein A
Specificity	This antibody does not cross-react with human HMGN2, HMGN3, HMGN4, HMGN5.
Concentration	0.531 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/200
Notes	

	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Transfected cell line - Human
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Rat, Transfected cell line - Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Rat, Transfected cell line - Human
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat, Transfected cell line - Human
Dilution info	-
Notes	-

WB

Not recommended

Species	Human, Mouse, Rat, Transfected cell line - Human
Dilution info	-
Notes	-

ChIP-seq

Not recommended

Species	Human, Mouse, Rat, Transfected cell line - Human
Dilution info	-
Notes	-

Target data

See full target information HMGN1

Function	Binds to the inner side of the nucleosomal DNA thus altering the interaction between the DNA and the histone octamer. May be involved in the process which maintains transcribable genes in a unique chromatin conformation. Inhibits the phosphorylation of nucleosomal histones H3 and H2A by RPS6KA5/MSK1 and RPS6KA3/RSK2 (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	HMGN1 also known as high mobility group nucleosomal binding domain protein 1 is a non-histone chromosomal protein with a mass of approximately 11 kDa. It interacts with chromatin inside the nucleus and regulates the architecture of

	nucleosomes. HMGN1 expresses in various types of tissues with higher levels observed in actively proliferating cells. The protein modulates the structure of chromatin by binding to nucleosomes and affecting nucleosome spacing and histone modifications.
Biological function summary	This protein plays an important role in modulating gene expression and cellular response to DNA damage. It does not form part of a large complex but associates with nucleosomes to influence chromatin dynamics. HMGN1 can affect transcription by altering chromatin accessibility and stabilizer that facilitates repair processes in response to DNA damage. The protein can also interfere with specific transcription factors therefore reducing or enhancing the transcription of target genes.
Pathways	This protein fits into cellular pathways involving transcription regulation and DNA repair. HMGN1 modifies chromatin structure and therefore participates in transcription-related pathways. It is also involved in DNA repair pathways ensuring proper response to DNA damage. Within these systems HMGN1 interacts with proteins such as HMGA1 influencing transcription and ATM which is critical in the DNA damage response pathway.
Associated diseases and disorders	Irregular expression or dysfunction of this protein links to certain cancers and autoimmune disorders. HMGN1 influences the development and progression of diseases through its role in gene expression and chromatin modulation. Aberrations in HMGN1 expression can lead to improper regulation of related proteins such as p53 an important player in cancer progression and NF-kB involved in immune responses. This connection with p53 and NF-kB makes HMGN1 a target of interest in disease research and therapy development.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

6 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HMGN1 antibody [EPR26772-17] (ab314864)

HMGN1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HMGN1 antibody

Immunohistochemical analysis of paraffin-embedded (A) HEK-293T (human embryonic kidney epithelial cell) transfected with a HMGN1 expression vector containing a Myc-His tag. (B) HEK-293T transfected with empty vector containing a Myc-His tag. tissue labeling HMGN1 with ab314864 at 1/2000 (0.266 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on HEK-293T transfected with a Myc-His-tagged HMGN1 construct (image A). No staining on HEK-293T transfected with empty plasmid (image B). The section was incubated with ab314864 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HMGN1 antibody [EPR26772-17] (ab314864)

Immunohistochemical analysis of paraffin-embedded human breast tissue labeling HMGN1 with ab314864 at 1/200 (2.655 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human breast. The section was incubated with ab314864 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HMGN1 antibody [EPR26772-17] (ab314864)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue labeling HMGN1 with ab314864 at 1/200 (2.655 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human breast carcinoma. The section was incubated with ab314864 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-HMGN1 antibody [EPR26772-17] (ab314864)


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NCCIT (human pluripotent embryonic carcinoma epithelial cell) cells labelling HMGN1 with ab314864 at 1/50 (10.34 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/mL) dilution (Green). Confocal image showing nuclear staining in NCCIT cell line. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/mL) dilution.



Flow Cytometry (Intracellular) - Anti-HMGN1 antibody [EPR26772-17] (ab314864)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NCCIT (human pluripotent embryonic carcinoma epithelial cell) cells labelling HMGN1 with ab314864 at 1/50 dilution (1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-HMGN1 antibody [EPR26772-17] (ab314864)

HMGN1 was immunoprecipitated from 0.35 mg NCCIT (human pluripotent embryonic carcinoma epithelial cell) whole cell lysate with ab314864 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab314864 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: NCCIT (human pluripotent embryonic carcinoma epithelial cell) whole cell lysate

Lane 2: ab314864 IP in NCCIT (human pluripotent embryonic carcinoma epithelial cell) whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab314864 in NCCIT whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-HMGN1 antibody [EPR26772-17] (ab314864) at 1/30 dilution

All lanes:
NCCIT (human pluripotent embryonic carcinoma epithelial cell) whole cell lysate

Secondary

All lanes:

Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.