abcam

ab314870

Anti-CLEC5A antibody [EPR28104-14]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR28104-14
Purification technique	Affinity purification Protein A
Concentration	0.528 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

Dot

Tested

Species	Recombinant fragment - Mouse
Dilution info	1/1000
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human, Rat

Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat, Recombinant fragment - Mouse
Dilution info	-
Notes	-

WB

Not recommended

Species	Human, Mouse, Rat, Recombinant fragment - Mouse
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

Target data

See full target information Clec5a

Function	Functions as a positive regulator of osteoclastogenesis (PubMed:19251634). Cell surface receptor that signals via TYROBP (PubMed:10449773). Regulates inflammatory responses (PubMed:20212065).

Storage

Shipped at conditions	Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.


Full details and terms and conditions can be found here:
Terms & Conditions.

5 product images




Immunocytochemistry/ Immunofluorescence - Anti-CLEC5A antibody [EPR28104-14] (ab314870)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) cells labelling CLEC5A with ab314870 at 1/50 (10.56 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/mL dilution (Green). Confocal image showing membrane and cytoplasmic staining in RAW 264.7 cell line. Negative control: EL4. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution.




Flow Cytometry - Anti-CLEC5A antibody [EPR28104-14] (ab314870)

Flow cytometric analysis of EL4 (mouse lymphoma T lymphocyte, Left) / RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage, Right) cells labelling CLEC5A with ab314870 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Negative control: EL4. Gated on viable cells.




Flow Cytometry - Anti-CLEC5A antibody [EPR28104-14] (ab314870)

Flow cytometric analysis of Mouse splenocytes cells labelling CLEC5A with ab314870 at 1/500 dilution (0.1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Gated on viable cells. Cells are co-stained with CD11b conjugated to Brilliant Violet™ 421.



Flow Cytometry - Anti-CLEC5A antibody [EPR28104-14] (ab314870)

Flow cytometric analysis of Mouse splenocytes cells labelling CLEC5A with ab314870 at 1/500 dilution (0.1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Gated on viable cells. Cells are co-stained with CD3 conjugated to Alexa Fluor®647.



Dot Blot - Anti-CLEC5A antibody [EPR28104-14] (ab314870)

Dot blot analysis of CLEC5A using ab314870 at 1:1000 (0.528 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1:100,000 dilution. Lane 1: His-tagged mouse CLEC5A fragment Exposure time: 180 seconds Blocking and diluting buffer and concentration: 5% NFDM/TBST

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.