abcam

ab314929

Alexa Fluor® 488 Anti-EPCR/CD201 antibody [EPR26267-19]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR26267-19
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/5000
Notes	-

Target data

See full target information Procr

Function	Binds activated protein C. Enhances protein C activation by the thrombin-thrombomodulin complex; plays a role in the protein C pathway controlling blood coagulation.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	EPCR also known as CD201 stands for Endothelial Protein C Receptor. It is a glycoprotein with a molecular mass of approximately 46 kDa. EPCR is expressed mainly on endothelial cells which line the interior surface of blood vessels. This receptor plays a mechanical role in binding to protein C a zymogen that is converted to activated protein C (APC) by thrombin-thrombomodulin complex. By facilitating the activation of protein C EPCR contributes to the regulation of blood coagulation and inflammation processes.
Biological function summary	The EPCR is part of the protein C anticoagulant pathway where it acts as a cofactor. EPCR is vital in enhancing the activation of protein C on the endothelial surface reducing coagulation activity. It interacts closely with thrombomodulin and protein C to form this complex preventing excessive thrombus formation and contributing to vascular integrity. Apart from its anticoagulant role EPCR also modulates inflammatory responses supporting its function as a bridge between coagulation and inflammation.
Pathways	EPCR lies within the protein C anticoagulation pathway and the endothelial signaling pathways. Its primary partner in the anticoagulation pathway is APC which degrades activated factors Va and VIIIa providing a check against coagulation. Additionally EPCR's interaction with APC and thrombomodulin links this protein to the endothelial signaling pathway where it affects cell survival and inflammation regulation. This interaction is essential to maintain hemostatic balance and control vascular responses to inflammatory stimuli.

Associated diseases and disorders	EPCR associates with conditions like thrombosis and sepsis. In thrombosis inadequate expression or function of EPCR can lead to uncontrolled blood clotting due to insufficient activation of protein C. Sepsis a severe inflammatory response to infection shows abnormal modulation of the protein C pathway where EPCR's role becomes pivotal. Additionally APC which interacts directly with EPCR is critical in the pathological states of these diseases highlighting the therapeutic potential in targeting EPCR pathway.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.


Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-EPCR/CD201 antibody [EPR26267-19] (ab314929)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized bEnd.3 (mouse brain endothelial cell) cells labelling EPCR/CD201 with ab314929 at 1/50 (10.0 ug/ml) dilution. Confocal image showing cytoplasmic staining in bEnd.3 cell line. Low expression: B16-F10 (PMID: 19188668). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue).



Flow Cytometry - Alexa Fluor® 488 Anti-EPCR/CD201 antibody [EPR26267-19] (ab314929)

Flow cytometric analysis of B16-F10 (mouse skin melanoma cell, Left) / bEnd.3 (mouse brain endothelial cell, Right) cells labelling EPCR/CD201 with ab314929 at 1/5000 dilution (0.01 ug)/Red compared with a Rabbit IgG monoclonal [EPR25A] - Isotype Control (Alexa Fluor® 488) (ab199091) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Low expression: B16-F10. Gated on viable cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.