abcam

ab315096

Anti-Cytokeratin 18 antibody [RM1051]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Multiclonal
Immunogens	The exact immunogen used to generate this antibody is proprietary information.
Clone number	RM1051
Purification technique	Affinity purification Protein A
Specificity	Unsuitable for Rat ICC.
Concentration	0.474 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Species	Mouse
Dilution info	1/50

Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Species	Mouse
Dilution info	1/500
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Species	Mouse
Dilution info	1/30

Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information KRT18

Function	Required for the formation of KRT8/KRT18 filaments that are involved in ARHGEF40-mediated actin stress fiber formation and tensional force-induced stress fiber formation and reinforcement (PubMed:26823019). Also acts downstream of ROCK kinase activation as part of a positive feedback mechanism in response to cellular mechanical stress loading (PubMed:26823019). Organization and orientation of KRT18 filaments are responsible for the properly elongated morphology of epithelial tubules (By similarity). Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What are recombinant multiclonals?
Recombinant multiclonals are a mixture of recombinant antibodies co-expressed from a library of heavy and light chains. They offer several advantages including:

- The sensitivity of polyclonal antibodies by recognising multiple epitopes
- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

View our range of recombinant multiclonal antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Cytokeratin 18 (CK18) also known as Keratin 18 or KRT18 is a type I intermediate filament protein with a molecular weight of approximately 45 kDa. It plays a mechanical role in stabilizing epithelial cell structure and function. CK18 is expressed in single-layered epithelial tissues such as the liver intestine and pancreas. It partners with the type II keratin usually KRT8 to form a stable cytoskeletal network in these cells providing structural integrity and resilience.
Biological function summary	CK18 maintains cellular integrity and mediates resistance to mechanical and non-mechanical stresses in epithelial cells. It forms heterodimers with KRT8 participating in a complex network of filaments important for maintaining cell shape and stability. This organization plays a role in managing cell survival during stress and apoptosis helping regulate cellular functions and responses in tissues where it is expressed.
Pathways	Cytokeratin 18 is involved in the apoptosis and cellular stress response pathways. It interacts with proteins such as caspases during apoptosis facilitating cytoskeletal reorganization and cell fragmentation. CK18 in coordination with KRT8 also participates in signaling pathways that regulate cell cycle and apoptotic processes underlining its importance in cellular turnover and tissue homeostasis.
Associated diseases and disorders	Elevated levels of CK18 fragments are often observed in cases of non-alcoholic steatohepatitis (NASH) and breast cancer. It serves as a biomarker for liver injury due to its abundance in hepatocytes and release upon cell death. In breast cancer alterations in CK18 expression can reflect disease progression where it interacts with other proteins such as vimentin indicating epithelial-to-mesenchymal transition a process that facilitates metastasis.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

18 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)


Immunohistochemical analysis of paraffin-embedded Human gastric cancer tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on human gastric cancer. The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Cytokeratin 18 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human cerebrum using rabbit Anti-Cytokeratin 18 antibody


Immunohistochemical analysis of paraffin-embedded Human cerebrum tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Negative control: no staining on human cerebrum. The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Cytokeratin 18 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human skeletal muscle using rabbit Anti-Cytokeratin 18 antibody


Immunohistochemical analysis of paraffin-embedded Human skeletal muscle tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Negative control: no staining on human skeletal muscle. The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)


Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on human colon.The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins

---




Flow Cytometry (Intracellular) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Molt-4 (human lymphoblastic leukemia T lymphoblast, Left) / A431 (human epidermoid carcinoma epithelial cell, Right) cells labelling Cytokeratin 18 with ab315096 at 1/500 dilution (0.1 ug, Red)

compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Negative control: Molt-4.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Immunohistochemical analysis of paraffin-embedded Human ovarian cancer tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on human ovarian cancer.The section was incubated with ab315096 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized A431 (human epidermoid carcinoma epithelial cell) cells labelling Cytokeratin 18 with ab315096 at 1/50 (9.48 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing cytoplasmic staining in A431 cell line. Negative control: MOLT-4.Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab7291 Anti-alpha Tubulin mouse monoclonal antibody was used to counterstain tubulin at 1/1000 (1 ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Immunohistochemical analysis of paraffin-embedded Human stomach tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on human stomach. The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins



Immunoprecipitation - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Cytokeratin 18 was immunoprecipitated from 0.35 mg A431 (human epidermoid carcinoma epithelial cell) whole cell lysate with ab315096 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab315096 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: A431 (human epidermoid carcinoma epithelial cell) whole cell lysate

Lane 2: ab315096 IP in A431 (human epidermoid carcinoma epithelial cell) whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab315096 in A431 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


All lanes:
Immunoprecipitation - Anti-Cytokeratin 18 antibody [RM1051] (ab315096) at 1/30 dilution

All lanes:
A431 (human epidermoid carcinoma epithelial cell) whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 15s




Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Hepa1-6 (mouse hepatoma epithelial cell) cells labelling Cytokeratin 18 with ab315096 at 1/50 (9.48 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing cytoplasmic staining in Hepa1-6 cell line. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab7291 Anti-alpha Tubulin mouse monoclonal antibody was used to counterstain tubulin at 1/1000 (1 ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Immunohistochemical analysis of paraffin-embedded Mouse stomach tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on mouse stomach. The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Immunohistochemical analysis of paraffin-embedded Rat stomach tissue labeling Cytokeratin 18 with ab315096 at 1/500 (0.948 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on rat stomach. The section was incubated with ab315096 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins



Flow Cytometry (Intracellular) - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Hepa1-6 (mouse hepatoma epithelial cell) cells labelling Cytokeratin 18 with ab315096 at 1/500 dilution (0.1 ug, Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Cytokeratin 18 was immunoprecipitated from 0.35 mg Hepa1-6 (mouse hepatoma epithelial cell) whole cell lysate with ab315096 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab315096 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: Hepa1-6 (mouse hepatoma epithelial cell) whole cell lysate

Lane 2: ab315096 IP in Hepa1-6 (mouse hepatoma epithelial cell) whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab315096 in Hepa1-6 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


All lanes:
Immunoprecipitation - Anti-Cytokeratin 18 antibody [RM1051] (ab315096) at 1/30 dilution

All lanes:
Hepa1-6 (mouse hepatoma epithelial cell) whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 15s



Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Low expression: Skeletal muscle.

Generation of the 22kDa fragment can be induced by staurosporine treatment (lane 1-2) (PMID: 34322741 and PMID:9298992).

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

Exposure time: Lanes 1-2: 8 seconds; Lanes 3-5: 1 second.

All lanes:
Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096) at 1/1000 dilution

Lane 1:
Untreated HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
HeLa treated with 2uM staurosporine for 24 hours whole cell lysate at 20 µg

Lane 3:
Human lung tissue lysate at 20 µg


Lane 4:
Human liver tissue lysate at 20 µg

Lane 5:
Human skeletal muscle tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 22 kDa, 48 kDa, 36 kDa



Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Negative control: MOLT-4 (PMID: 2473821)

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096) at 1/1000 dilution

Lane 1:
A431 (human epidermoid carcinoma epithelial cell) whole cell lysate at 7 µg


Lane 2:
MOLT-4 (human lymphoblastic leukemia T lymphoblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 48 kDa, 36 kDa

Exposure time: 4s



Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Cytokeratin 18 Western blot staining using rabbit Anti-Cytokeratin 18 antibody

Western blot: Anti-Cytokeratin 18 antibody [RM1051] ab315096 staining at 1/1000 dilution, shown in green; Mouse anti GAPDH ab8245 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 48 kDa in Wild-type A549 cell line with no signal observed at this size in KRT18 knockout A549 cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5% Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096) at 1/1000 dilution

Lane 1:
Wild-type A549 cell lysates at 20 µg

Lanes 2 and 4:
Empty

Lane 3:
KRT18 knockout A549 cell lysates at 20 µg

Lane 5:
HeLa cell lysates at 20 µg

Lane 6:
SH-SY5Y cell lysates at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 48 kDa

Observed band size: 48 kDa



Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096)

Cytokeratin 18 Western blot staining using rabbit Anti-Cytokeratin 18 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Low expression: Skeletal muscle.

The observed MW is consistent with what has been described in the literature (PMID:9298992).

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

Exposure time: Lanes 1: 26 seconds; Lanes 2-7: 180 seconds.

All lanes:
Western blot - Anti-Cytokeratin 18 antibody [RM1051] (ab315096) at 1/1000 dilution

Lane 1:
Hepa1-6 (mouse hepatoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Mouse colon tissue lysate at 20 µg

Lane 3:
Mouse liver tissue lysate at 20 µg

Lane 4:
Mouse skeletal muscle tissue lysate at 20 µg

Lane 5:
Rat liver tissue lysate at 20 µg

Lane 6:
Rat skeletal muscle tissue lysate at 20 µg

Lane 7:
AR42J (rat pancreatic tumor epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 45 kDa, 48 kDa, 36 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.