abcam

ab315971

Anti-Pofut1 antibody [EPR26782-194]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR26782-194
Purification technique	Affinity purification Protein A
Concentration	0.514 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse

Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/1000
Notes	

	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

IP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information POFUT1

Function	Catalyzes the reaction that attaches fucose through an O-glycosidic linkage to a conserved serine or threonine residue found in the consensus sequence C2-X(4,5)-[S/T]-C3 of EGF domains, where C2 and C3 are the second and third conserved cysteines. Specifically uses GDP-fucose as donor substrate and proper disulfide pairing of the substrate EGF domains is required for fucose transfer. Plays a crucial role in NOTCH signaling. Initial fucosylation of NOTCH by POFUT1 generates a substrate for FRINGE/RFNG, an acetylglucosaminyltransferase that can then extend the fucosylation on the NOTCH EGF repeats. This extended fucosylation is required for optimal ligand binding and canonical NOTCH signaling induced by DLL1 or JAGGED1. Fucosylates AGRN and determines its ability to cluster acetylcholine receptors (AChRs).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The protein known as Pofut1 or protein O-fucosyltransferase 1 functions mechanically by adding fucose a sugar molecule to serine or threonine residues on proteins specifically within the epidermal growth factor-like (EGF-like) repeats of substrates. Pofut1 has a molecular weight of approximately 50 kDa. This protein is expressed in various tissues including heart liver and brain indicating its significant role across different biological systems. Researchers commonly refer to Pofut1 in studies focused on protein glycosylation processes.
Biological function summary	Pofut1 is responsible for the proper folding and functioning of specific proteins that undergo fucosylation. It does not technically form part of a multi-protein complex but interacts tightly with substrates it modifies. Through its enzymatic action Pofut1 influences essential biological activities in cell communication and signaling. This glycosyltransferase also plays a pivotal role in modulating Notch signaling a critical pathway in the development and tissue regeneration processes.

Pathways	The significance of Pofut1 is most notable within the Notch signaling pathway where it modifies the Notch receptor enabling effective ligand-receptor interactions. Dysregulation of this pathway can lead to critical developmental defects. Pofut1 also connects to other pathways such as the Fringe-mediated glycosylation pathway. In this context Pofut1-related proteins like Fringe glycosyltransferases operate to further influence Notch signaling outcomes through additional glycosylation processing.
Associated diseases and disorders	The functionality of Pofut1 has links to specific syndromes and malignancies. For instance aberrations in its activity are associated with congenital heart defects due to disrupted Notch signaling affecting cardiac development. Additionally overexpression or mutations in Pofut1 relate to cancer as improper Notch signaling can lead to tumor progression. Relatedly proteins like Notch1 serve as intermediaries linking Pofut1 activity to such conditions thereby highlighting Pofut1's involvement in disease etiologies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


10 product images



Flow Cytometry (Intracellular) - Anti-Pofut1 antibody [EPR26782-194] (ab315971)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HT-29 (human colorectal adenocarcinoma epithelial cell) cells labelling Pofut1 with ab315971 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Pofut1 antibody [EPR26782-194] (ab315971)


Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling Pofut1 with ab315971 at 1/1000 (0.514 ug/ml) dilution, followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Positive staining on human kidney.
The section was incubated with ab315971 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Pofut1 antibody [EPR26782-194] (ab315971)


Immunohistochemical analysis of paraffin-embedded Human placenta tissue labeling Pofut1 with ab315971 at 1/1000 (0.514 ug/ml) dilution, followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Positive staining on human placenta (PMID: 11524432; PMID: 25165882).
The section was incubated with ab315971 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Pofut1 antibody [EPR26782-194] (ab315971)


Immunohistochemical analysis of paraffin-embedded Rat kidney tissue labeling Pofut1 with ab315971 at 1/1000 (0.514 ug/ml) dilution, followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Positive staining on rat kidney.
The section was incubated with ab315971 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Pofut1 antibody [EPR26782-194] (ab315971)


Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labeling Pofut1 with ab315971 at 1/1000 (0.514 ug/ml) dilution, followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Positive staining on mouse kidney (PMID: 12697902).
The section was incubated with ab315971 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-Pofut1 antibody [EPR26782-194] (ab315971)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized C2C12 (mouse myoblast) cells labelling Pofut1 with ab315971 at 1/50 dilution (1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971)

Pofut1 Western blot staining using rabbit Anti-Pofut1 antibody

All lanes:
Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971) at 1/1000 dilution

Lane 1:
HT-29 (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
MOLT-4 (human lymphoblastic leukemia T lymphoblast) whole cell lysate at 20 µg

Lane 3:
THP-1 (human monocytic leukemia monocyte) whole cell lysate at 20 µg

Lane 4:
C2C12 (mouse myoblast) whole cell lysate at 20 µg

Lane 5:
bEnd.3 (mouse brain endothelial cell) whole cell lysate at 20 µg

Lane 6:
PC-12 (rat adrenal gland pheochromocytoma cell) whole cell lysate at 20 µg


Lane 7:
C6 (rat glial tumor glial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 44 kDa

Exposure time: 26s



Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971)

Pofut1 Western blot staining using rabbit Anti-Pofut1 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The identity of the lower MW band at approximately 75kDa in Lane 3 is unknown.

All lanes:
Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971) at 1/1000 dilution

Lane 1:
Human liver tissue lysate at 20 µg

Lane 2:
Mouse liver tissue lysate at 20 µg


Lane 3:
Rat kidney tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 44 kDa

Exposure time: 180s



Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971)

Pofut1 Western blot staining using rabbit Anti-Pofut1 antibody

Western blot: Anti-Pofut1 antibody [EPR26782-194] ab315971 staining at 1/1000 dilution, shown in green; Mouse anti-CANX ab238078 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 44 kDa in Wild-type A549 cell line with no signal observed at this size in POFUT1 knockout A549 cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5% Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971) at 1/1000 dilution

Lane 1:
Wild-type A549 cell lysates at 20 µg

Lane 2:
POFUT1 knockout A549 cell lysates at 20 µg

Lane 3:
HCT 116 cell lysates at 20 µg

Lane 4:
THP-1 cell lysates at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 44 kDa

Observed band size: 44 kDa



Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971)

Pofut1 Western blot staining using rabbit Anti-Pofut1 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

In Western blot, Anti-Vinculin antibody [EPR8185] (ab129002) staining at 1/10000 dilution.

All lanes:
Western blot - Anti-Pofut1 antibody [EPR26782-194] (ab315971) at 1/1000 dilution

Lane 1:
HCT 116 (human colorectal carcinoma epithelial cell) transfected with scrambled siRNA control whole cell lysate at 20 µg

Lane 2:
HCT 116 transfected with siRNA specifically targeting Pofut1 whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 44 kDa, 124 kDa

Exposure time: 26s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.