abcam

ab316017

Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR28780-7
Purification technique	Affinity purification Protein A
Concentration	1.028 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Dot

Tested

Species	Recombinant full length protein - Human
Dilution info	-

Notes	-

Expected

Species	Human, Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat, Recombinant full length protein - Human
Dilution info	-
Notes	-

Target data

See full target information AKR1B1

Function	Catalyzes the NADPH-dependent reduction of a wide variety of carbonyl-containing compounds to their corresponding alcohols. Displays enzymatic activity towards endogenous metabolites such as aromatic and aliphatic aldehydes, ketones, monosacharides, bile acids and xenobiotics substrates. Key enzyme in the polyol pathway, catalyzes reduction of glucose to sorbitol during hyperglycemia (PubMed:1936586). Reduces steroids and their derivatives and prostaglandins. Displays low enzymatic activity toward all-trans-retinal, 9-cis-retinal, and 13-cis-retinal (PubMed:12732097, PubMed:19010934, PubMed:8343525). Catalyzes the reduction of diverse phospholipid aldehydes such as 1-palmitoyl-2-(5-oxovaleroyl)-sn -glycero-3-phosphoethanolamin (POVPC) and related phospholipid aldehydes that are generated from the oxydation of phosphotidylcholine and phosphatdyleethanolamides (PubMed:17381426). Plays a role in detoxifying dietary and lipid-derived unsaturated carbonyls, such as crotonaldehyde, 4-hydroxynonenal, trans-2-hexenal, trans-2,4-hexadienal and their glutathione-conjugates carbonyls (GS-carbonyls) (PubMed:21329684).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

ab316017 is the carrier-free version of ab316016.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Aldose reductase also known as AR or ALR2 is an important enzyme in the polyol pathway. It has a molecular mass of about 36 kDa and reduces aldehydes including glucose to their corresponding alcohols. Aldose reductase is expressed widely in various tissues such as kidney lens and retina. The enzyme uses NADPH as a cofactor which is important for its catalytic activity.
Biological function summary	The main function of aldose reductase involves converting glucose to sorbitol a process particularly active under hyperglycemic conditions. It does not form a part of a larger complex but works individually to facilitate this conversion. Sorbitol the product of its enzymatic reaction eventually converts to fructose through the action of sorbitol dehydrogenase. This two-step conversion can contribute to cellular damage due to osmotic stress when sorbitol accumulates.
Pathways	Aldose reductase plays a central role in the polyol pathway. This pathway becomes highly relevant when blood glucose levels are elevated. During such conditions aldose reductase activity increases to convert excess glucose to sorbitol. This activity links it to other proteins like sorbitol dehydrogenase which completes the conversion to fructose.
Associated diseases and disorders	Aldose reductase has strong connections to diabetic complications specifically diabetic neuropathy and retinopathy. The enzyme's increased activity under high glucose conditions can lead to osmotic stress and tissue damage. Interactions with other proteins in the metabolic shift associated with diabetes such as sorbitol dehydrogenase underline its involvement. Aldose reductase inhibitors show potential in mitigating these adverse effects by reducing sorbitol accumulation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

16 product images




Flow Cytometry (Intracellular) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HeLa (human cervical adenocarcinoma epithelial cell) cells labelling Aldose reductase with ab316016 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)


This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HeLa (human cervical adenocarcinoma epithelial cell) cells labelling Aldose reductase with ab316016 at 1/100 (5.09 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/mL) dilution (Green).

Confocal image showing cytoplasmic and nuclear staining in HeLa cell line.
Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/mL) dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

This data was developed using ab316016, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Human adrenal gland tissue labeling Aldose reductase with ab316016 at 1/6000 (0.085 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on human adrenal gland (PMID: 32602410).
The section was incubated with ab316016 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human liver using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling Aldose reductase with ab316016 at 1/6000 (0.085 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Negative control: No staining on human liver.
The section was incubated with ab316016 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

This data was developed using ab316016, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Mouse adrenal gland tissue labeling Aldose reductase with ab316016 at 1/6000 (0.085 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on mouse adrenal gland.
The section was incubated with ab316016 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse liver using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Mouse liver tissue labeling Aldose reductase with ab316016 at 1/6000 (0.085 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Negative control: No staining on mouse liver.
The section was incubated with ab316016 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Rat liver using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Rat liver tissue labeling Aldose reductase with ab316016 at 1/6000 (0.085 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Negative control: No staining on rat liver.
The section was incubated with ab316016 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

This data was developed using ab316016, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Rat adrenal gland tissue labeling Aldose reductase with ab316016 at 1/6000 (0.085 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on rat adrenal gland.
The section was incubated with ab316016 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)


This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized PC-12 (rat adrenal gland pheochromocytoma cell) cells labelling Aldose reductase with ab316016 at 1/100 (5.09 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/mL) dilution (Green).

Confocal image showing cytoplasmic and nuclear staining in PC-12 cell line.
Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/mL) dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)


This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling Aldose reductase with ab316016 at 1/100 (5.09 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/mL) dilution (Green).

Confocal image showing cytoplasmic and nuclear staining in NIH/3T3 cell line.
Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/mL) dilution.




Flow Cytometry (Intracellular) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling Aldose reductase with ab316016 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.




Flow Cytometry (Intracellular) - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized PC-12 (rat adrenal gland pheochromocytoma cell) cells labelling Aldose reductase with ab316016 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Western blot - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Western blot staining using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-Aldose reductase antibody [EPR28780-7] (ab316016) at 1/1000 dilution

Lane 1:
HeLa (human cervical adenocarcinoma epithelial cell) transfected with scrambled siRNA control whole cell lysate at 20 µg

Lane 2:
HeLa transfected with siRNA specifically targeting Aldose reductase whole cell lysate at 20 µg

Lane 3:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 4:
PC-12 (rat adrenal gland pheochromocytoma cell) whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Developed using the ECL technique.

Observed band size: 36 kDa

Exposure time: 180s



Western blot - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Western blot staining using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Negative control: liver

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-Aldose reductase antibody [EPR28780-7] (ab316016) at 1/1000 dilution

Lane 1:
Human adrenal gland tissue lysate at 20 µg

Lane 2:
Human testis tissue lysate at 20 µg


Lane 3:
Human liver tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 36 kDa

Exposure time: 81s



Western blot - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Western blot staining using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Negative control: liver

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-Aldose reductase antibody [EPR28780-7] (ab316016) at 1/1000 dilution

Lane 1:
Mouse adrenal gland tissue lysate at 20 µg

Lane 2:
Mouse testis tissue lysate at 20 µg

Lane 3:
Rat adrenal gland tissue lysate at 20 µg

Lane 4:
Rat testis tissue lysate at 20 µg


Lane 5:
Rat liver tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 36 kDa

Exposure time: 81s



Dot Blot - Anti-Aldose reductase antibody [EPR28780-7] - BSA and Azide free (ab316017)

Aldose reductase Dot Blot staining using rabbit Anti-Aldose reductase antibody

This data was developed using ab316016, the same antibody clone in a different buffer formulation.

Dot blot analysis of Aldose reductase using ab316016 at 1:1000 (0.509 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1:100,000 dilution.

This antibody does not cross-react with human AKR1B10 and AKR1B15.

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

All lanes:
Dot Blot - Anti-Aldose reductase antibody [EPR28780-7] (ab316016) at 1/1000 dilution

Lane 1:
His-tagged human Aldose reductase full length recombinant protein

Lane 2:
His-tagged human AKR1B10 full length recombinant protein

Lane 3:
His-tagged human AKR1B15 full length recombinant protein

Secondary

All lanes:
Dot Blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.