abcam

ab316306

Anti-CBLL1 antibody [EPR25389-121]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR25389-121
Purification technique	Affinity purification Protein A
Concentration	0.513 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse

Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Species	Mouse
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/2000
Notes	

	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	-

Species	Mouse
Dilution info	1/100
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Species	Mouse
Dilution info	1/500
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information CBLL1

Function	E3 ubiquitin-protein ligase that mediates ubiquitination of several tyrosine-phosphorylated Src substrates, including CDH1, CTTN and DOK1 (By similarity). Targets CDH1 for endocytosis and degradation (By similarity). Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing (PubMed:29507755). Its function in the WMM complex is unknown (PubMed:29507755).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CBLL1 also known as Hakai is an E3 ubiquitin ligase. Its molecular weight is approximately 65 kDa and it functions by mediating ubiquitination which leads to the controlled degradation of target proteins. CBLL1 is expressed in several tissues including heart liver and kidney. The protein predominantly functions within the cytoplasm but can also translocate to the nucleus to exert its enzymatic activities.
Biological function summary	CBLL1 plays a significant role in cellular processes such as endocytosis cell adhesion and regulation of epithelial-mesenchymal transition (EMT). CBLL1 functions as part of a complex that helps target proteins for degradation through the ubiquitin-proteasome system. This activity is especially relevant in the regulation of cell signaling pathways where it targets various substrates such as E-cadherin for ubiquitination.
Pathways	CBLL1 is important in modulating signaling cascades like the Wnt and TGF-beta pathways. By influencing these pathways CBLL1 interacts with proteins like beta-catenin and SMADs. The degradation of E-cadherin by CBLL1 facilitates cellular detachment and migration a critical step in cellular processes regulated by these pathways.
Associated diseases and disorders	CBLL1's involvement in the ubiquitin-proteasome system links it to cancer development and progression. The protein's regulation of EMT and cell adhesion implicates it in cancers such as gastric and colon cancers. Additionally CBLL1 interacts with proteins like beta-catenin in Wnt signaling which is often dysregulated in these cancers. Understanding the function of CBLL1 may offer potential therapeutic targets for combating these malignancies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


13 product images



Flow Cytometry (Intracellular) - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized MCF7 (human breast adenocarcinoma epithelial cell) cells labelling CBLL1 with ab316306 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling CBLL1 with ab316306 at 1/1000 (0.513 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Positive staining on human tonsil (PMID: 19535458). The section was incubated with ab316306 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CBLL1 antibody [EPR25389-121] (ab316306)


Immunohistochemical analysis of paraffin-embedded Human gastric carcinoma tissue labeling CBLL1 with ab316306 at 1/1000 (0.513 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on human gastric carcinoma. The section was incubated with ab316306 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized MCF7 (human breast adenocarcinoma epithelial cell) cells labelling CBLL1 with ab316306 at 1/100 (5.13 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing mainly nuclear staining in MCF7 cell line.
Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Immunoprecipitation - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

CBLL1 was immunoprecipitated from 0.35 mg MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate with ab316306 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab316306 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate
Lane 2: ab316306 IP in MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab316306 in MCF7 whole cell lysate


All lanes:
Immunoprecipitation - Anti-CBLL1 antibody [EPR25389-121] (ab316306) at 1/30 dilution

All lanes:
MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 180s



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CBLL1 antibody [EPR25389-121] (ab316306)


Immunohistochemical analysis of paraffin-embedded Mouse lung tissue labeling CBLL1 with ab316306 at 1/2000 (1.026 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on mouse lung. The section was incubated with ab316306 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling CBLL1 with ab316306 at 1/100 (5.13 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing nuclear staining in NIH/3T3 cell line.
Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Flow Cytometry (Intracellular) - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling CBLL1 with ab316306 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CBLL1 antibody [EPR25389-121] (ab316306)


Immunohistochemical analysis of paraffin-embedded Rat liver tissue labeling CBLL1 with ab316306 at 1/2000 (1.026 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on rat liver. The section was incubated with ab316306 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins



Immunoprecipitation - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

CBLL1 was immunoprecipitated from 0.35 mg NIH/3T3 (mouse embryonic fibroblast) whole cell lysate with ab316306 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab316306 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP) (ab131366) was used at 1/5000 dilution.

Lane 1: NIH/3T3 (mouse embryonic fibroblast) whole cell lysate
Lane 2: ab316306 IP in NIH/3T3 (mouse embryonic fibroblast) whole cell lysate
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab316306 in NIH/3T3 whole cell lysate


All lanes:
Immunoprecipitation - Anti-CBLL1 antibody [EPR25389-121] (ab316306) at 1/30 dilution

All lanes:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 24s



Western blot - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

CBLL1 Western blot staining using rabbit Anti-CBLL1 antibody

Lysates were freshly made and used for Western blotting immediately to minimize protein degradation.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-CBLL1 antibody [EPR25389-121] (ab316306) at 1/1000 dilution

Lane 1:
MCF7 (human breast adenocarcinoma epithelial cell) whole cell fresh lysate at 20 µg

Lane 2:
NIH/3T3 (mouse embryonic fibroblast) whole cell fresh lysate at 20 µg


Lane 3:
PC-12 (rat adrenal gland pheochromocytoma cell) whole cell fresh lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 55-60 kDa, 36 kDa

Exposure time: 37s



Western blot - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

CBLL1 Western blot staining using rabbit Anti-CBLL1 antibody

Lysates were freshly made and used for Western blotting immediately to minimize protein degradation.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-CBLL1 antibody [EPR25389-121] (ab316306) at 1/1000 dilution

Lane 1:
HeLa (human cervix adenocarcinoma epithelial cell) transfected with scrambled siRNA control whole cell lysate at 20 µg


Lane 2:
HeLa transfected with siRNA specifically targeting human CBLL1 whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 55-60 kDa, 36 kDa

Exposure time: 48s



Western blot - Anti-CBLL1 antibody [EPR25389-121] (ab316306)

CBLL1 Western blot staining using rabbit Anti-CBLL1 antibody

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-CBLL1 antibody [EPR25389-121] (ab316306) at 1/1000 dilution

Lane 1:
Mouse liver tissue lysate at 20 µg

Lane 2:
Mouse lung tissue lysate at 20 µg

Lane 3:
Rat liver tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 55-60 kDa, 36 kDa

Exposure time: 92s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.