abcam

ab317022

Anti-CD21+CD35 antibody [EPR27417-513]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR27417-513
Purification technique	Affinity purification Protein A
Concentration	0.499 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Dot

Tested

Species	Recombinant fragment - Mouse
Dilution info	1/1000
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/100
Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/5000
Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat

Dilution info	-
Notes	-

WB

Not recommended

Species	Human, Mouse, Rat, Recombinant fragment - Mouse
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat, Recombinant fragment - Mouse
Dilution info	-
Notes	-

Target data

See full target information Cr2

Function

Serves as a receptor for various ligands including complement component CD3d, HNRNPU OR IFNA1. When C3d is bound to antigens, attaches to C3d on B-cell surface and thereby facilitates the recognition and uptake of antigens by B-cells. This interaction enhances B-cell activation and subsequent immune responses. Forms a complex with several partners on the surface of B-cells including CD19, FCRL5 and CD81, to form the B-cell coreceptor complex that plays a crucial role in B-cell activation and signaling. Also induces specific intracellular signaling separately from the BCR and CD19 by activating the tyrosine kinase SRC, which then phosphorylates nucleolin/NCL and triggers AKT and GSK3 kinase activities in a SYK/CD19-independent manner. Acts as a ligand for CD23 (FcepsilonRII), a low-

affinity receptor for IgE, which is expressed on B-cells and other immune cells, and thus participates in the regulation of IgE production.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

5 product images




Immunocytochemistry/ Immunofluorescence - Anti-CD21+CD35 antibody [EPR27417-513] (ab317022)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Tween-20 permeabilized mouse splenocyte cells labelling CD21+CD35 with ab317022 at 1/100 (4.99 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing cytoplasmic staining in subsets of mouse splenocytes (shown in green). The counterstain was observed in magenta. Nuclear DNA was labeled with DAPI (shown in blue). Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Flow Cytometry - Anti-CD21+CD35 antibody [EPR27417-513] (ab317022)

Flow cytometric analysis of EL4 (mouse lymphoma T lymphocyte, Left) / WEHI-231 (mouse B cell lymphoma B lymphocyte, Right) cells labelling CD21+CD35 with ab317022 at 1/5000 dilution (0.01 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cell.
Negative control: EL4(PMID: 10453011)




Flow Cytometry - Anti-CD21+CD35 antibody [EPR27417-513] (ab317022)

CD21+CD35 Flow Cytometry staining of mouse splenocytes using rabbit Anti-CD21+CD35 antibody

Flow cytometric analysis of mouse splenocytes cells labelling CD21+CD35 with ab317022 at 1/5000 dilution (0.01 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

mouse splenocytes are co-stained with CD3 conjugated Alexa Fluor®647.
Gated on viable cell.




Flow Cytometry - Anti-CD21+CD35 antibody [EPR27417-513] (ab317022)

CD21+CD35 Flow Cytometry staining of mouse splenocytes using rabbit Anti-CD21+CD35 antibody

Flow cytometric analysis of mouse splenocytes cells labelling CD21+CD35 with ab317022 at 1/5000 dilution (0.01 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

mouse splenocytes are co-stained with CD19 conjugated Alexa Fluor®647.
Gated on viable cell.



Dot Blot - Anti-CD21+CD35 antibody [EPR27417-513] (ab317022)

CD21+CD35 Dot Blot staining of Mouse CR2 protein using rabbit Anti-CD21+CD35 antibody

Dot blot analysis of CD21+CD35 using ab317022 at 1:1000 (0.499 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1:100,000 dilution.

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

All lanes:
Dot Blot - Anti-CD21+CD35 antibody [EPR27417-513] (ab317022) at 1/1000 dilution

All lanes:
Mouse CR2 protein

Secondary

All lanes:
Dot Blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.