abcam

ab317272

Anti-LY6C antibody [RM1151]

Anti-LY6C antibody [RM1151] (ab317272) is a rabbit recombinant multiclonal antibody detecting Ly6c in **Western Blot, Flow Cytometry, IP, IHC-P, ICC/IF, mIHC**. Suitable for **Mouse**.

- Multiplex IHC validated on the Leica BOND® MAX using Opal reagents
- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Multiclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	RM1151
Purification technique	Affinity purification Protein A
Concentration	0.484 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

mIHC

Tested

Species	Mouse
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Human, Rat

Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30

Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Target data

See full target information Ly6c1

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-LY6C antibody [RM1151] (ab317272) is a rabbit recombinant recombinant multiclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Flow Cyt), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF), Multiplex IHC (mIHC) in Mouse samples.

Other related products
We have a range of other formats of antibody clone [RM1151] also available for your convenience: ab317272, Carrier free - ab317273

What are recombinant multiclonals?
Recombinant multiclonals are a mixture of recombinant antibodies co-expressed from a library of heavy and light chains. They offer several advantages including:

- The sensitivity of polyclonal antibodies by recognising multiple epitopes
- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

View our range of recombinant multiclonal antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Ly6c also known as Ly-6C is a protein marker with a molecular weight of approximately 18 kDa. It is part of the Ly6 family of glycosylphosphatidylinositol (GPI)-anchored cell surface proteins. Ly6c is predominantly expressed in mice on the surface of certain immune cells including monocytes macrophages dendritic cells and subsets of T cells. The expression levels can vary depending on the cell type and the activation or developmental stage of the cell. Ly6c serves as a useful marker in immunological studies to distinguish between different subsets of monocytes commonly identified as Ly6c^high and Ly6c^low populations.
Biological function summary	Ly6c is involved in the differentiation and trafficking of immune cells. It is not known to be part of a larger protein complex. In the immune system the Ly6c marker is critical for distinguishing between classical and non-classical monocytes in mice. Classical monocytes expressing high levels of Ly6c are involved in rapid response to inflammatory stimuli while non-classical monocytes with lower Ly6c expression help in patrolling blood vessels. This distinction is important in understanding how the immune system responds to various infections and how the body maintains immune homeostasis.
Pathways	Ly6c plays a role in the monocyte recruitment and differentiation pathways. It is closely related to the chemokine signaling pathway influencing how immune cells migrate towards sites of inflammation or injury. The presence of the Ly6c2 protein an isoform of Ly6c helps modulate these responses by interacting with cytokines and chemokines that guide the trafficking of immune cells. The presence of Ly6c influences the activity of proteins like CCR2 which plays a significant role in chemotaxis in response to inflammation.
Associated diseases and disorders	Ly6c is associated with inflammatory and autoimmune conditions. For example high Ly6c expression has been linked to inflammatory bowel disease (IBD) and atherosclerosis. In IBD the ability of Ly6c^high monocytes to migrate to the gut and promote inflammation is well-documented. Similarly in atherosclerosis the recruitment of Ly6c^high monocytes to blood vessels contributes to plaque formation. In these contexts proteins like MCP-1 (also known as CCL2) interact with

Ly6c-expressing cells facilitating their movement and function in disease processes. Understanding the role of Ly6c in these conditions might help develop targeted therapies to mitigate chronic inflammation and related diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images



Multiplex immunohistochemistry - Anti-LY6C antibody [RM1151] (ab317272)

LY6C Multiplex immunohistochemistry staining of Mouse spleen using rabbit Anti-LY6C antibody

Multiplex immunohistochemistry analysis of formalin/PFA-fixed paraffin-embedded mouse spleen tissue staining LY6C with ab317272 at a 1:100 (0.978 ug/ml) dilution, ab133357 anti-CD11b used at 1:20000 (0.05 ug/ml) dilution and ab245235 anti-CD19 used at a 1:1000 (0.999 ug/ml) dilution.


Panel A: merged staining of anti-Ly6c (green; Opal™690), anti-CD11b (magenta; Opal™570) and anti-CD19 (yellow; Opal™520) on mouse spleen.
Panel B: anti-Ly6c stained on monocytes/macrophages.
Panel C: anti-CD11b stained on monocytes/macrophages.
Panel D: anti-CD19 stained on B cells.
Co-staining of Ly6c and CD11b can be observed.
The section was incubated in three rounds of staining: in the order of ab317272, ab133357, and

ab245235 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.
Nuclear DNA was labeled with DAPI (shown in blue).

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.

Nuclear counter stain with DAPI.

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Flow Cytometry - Anti-LY6C antibody [RM1151] (ab317272)

Flow cytometric analysis of Mouse spleen cell cells labelling LY6C with ab317272 at 1/500 dilution (0.1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Mouse spleen cell are co-stained with CD8a conjugated Alexa Fluor®647.
Gated on viable cell.




Flow Cytometry - Anti-LY6C antibody [RM1151] (ab317272)

Flow cytometric analysis of Mouse bone marrow cells labelling LY6C with ab317272 at 1/500 dilution (0.1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Mouse bone marrow are co-stained with B220 conjugated Alexa Fluor®647.
Gated on viable cell.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LY6C antibody [RM1151] (ab317272)

LY6C Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse liver using rabbit Anti-LY6C antibody

Immunohistochemical analysis of paraffin-embedded Mouse liver tissue labeling LY6C with ab317272 at 1/500 (0.978 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Low expression tissue: no staining on mouse liver. The section was incubated with ab317272 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LY6C antibody [RM1151] (ab317272)

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labeling LY6C with ab317272 at 1/500 (0.978 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Positive staining on endothelial cells and immune cells in mouse colon. The section was incubated with ab317272 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LY6C antibody [RM1151] (ab317272)


Immunohistochemical analysis of paraffin-embedded Mouse cerebrum tissue labeling LY6C with ab317272 at 1/500 (0.978 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on endothelial cells in mouse cerebrum. The section was incubated with ab317272 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LY6C antibody [RM1151] (ab317272)


Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling LY6C with ab317272 at 1/500 (0.978 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on endothelial cells and immune cells in mouse spleen. The section was incubated with ab317272 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-LY6C antibody [RM1151] (ab317272)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse splenocyte cells labelling LY6C with ab317272 at 1/500 (0.978 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing cytoplasmic staining in subsets of mouse splenocytes (shown in green), which is partially co-located with CD11b. The counterstain was observed in magenta. Nuclear DNA was labeled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

ab8878 Anti-CD11b Rat monoclonal antibody was used to counterstain at 1/200 (2.5ug/ml) dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Immunocytochemistry/ Immunofluorescence - Anti-LY6C antibody [RM1151] (ab317272)


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse bone marrow cells labelling LY6C with ab317272 at 1/500 (0.978 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green).

Confocal image showing cytoplasmic staining in subsets of mouse bone marrow (shown in green), which is partially co-located with CD11b. The counterstain was observed in magenta. Nuclear DNA

was labeled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

ab8878 Anti-CD11b Rat monoclonal antibody was used to counterstain at 1/200 (2.5ug/ml) dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Flow Cytometry - Anti-LY6C antibody [RM1151] (ab317272)

Flow cytometric analysis of Mouse bone marrow cells labelling LY6C with ab317272 at 1/500 dilution (0.1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Mouse bone marrow are co-stained with CD11b conjugated BV510 (512/25BP).
Gated on viable cell.



Immunoprecipitation - Anti-LY6C antibody [RM1151] (ab317272)

LY6C was immunoprecipitated from 0.35 mg Mouse spleen tissue lysate with ab317272 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab317272 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: Mouse spleen tissue lysate
Lane 2: ab317272 IP in Mouse spleen tissue lysate
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab317272 in mouse spleen tissue lysate.


All lanes:
Immunoprecipitation - Anti-LY6C antibody [RM1151] (ab317272) at 1/30 dilution

All lanes:
Mouse spleen tissue lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 128s



Western blot - Anti-LY6C antibody [RM1151] (ab317272)

LY6C Western blot staining using rabbit Anti-LY6C antibody

Negative control: Neuro-2a.

Low expression: testis, liver.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:

Western blot - Anti-LY6C antibody [RM1151] (ab317272) at 1/1000 dilution

Lane 1:
Mouse lung tissue lysate at 20 µg

Lane 2:
Mouse spleen tissue lysate at 20 µg

Lane 3:
Mouse testis tissue lysate at 20 µg

Lane 4:
Mouse liver tissue lysate at 20 µg

Lane 5:
Mouse kidney tissue lysate at 20 µg

Lane 6:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 7:
C2C12 (mouse myoblast) whole cell lysate at 20 µg

Lane 8:
Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 14 kDa, 36 kDa

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.