abcam

ab317324

Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR29048-48
Purification technique	Affinity purification Protein A
Concentration	0.996 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	-

Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information MRC2

Function	May play a role as endocytotic lectin receptor displaying calcium-dependent lectin activity. Internalizes glycosylated ligands from the extracellular space for release in an endosomal compartment via clathrin-mediated endocytosis. May be involved in plasminogen activation system controlling the extracellular level of PLAUR/PLAU, and thus may regulate protease activity at the cell surface. May contribute to cellular uptake, remodeling and degradation of extracellular collagen matrices. May play a role during cancer progression as well as in other chronic tissue destructive diseases acting on collagen turnover. May participate in remodeling of extracellular matrix cooperating with the matrix metalloproteinases (MMPs).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	+4°C

Notes

ab317324 is the carrier-free version of ab317323.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	MRC2 also known as the Endo180 protein is a type 1 transmembrane receptor with a structure that includes a fibronectin type II domain eight C-type lectin domains and a cysteine-rich domain. This protein has an approximate molecular mass of 180 kDa. MRC2 is widely expressed in various human tissues with noticeable expression in fibroblasts macrophages and certain endothelial cells. The protein plays a role in endocytosis involving the binding and internalization of collagen and other extracellular matrix components.
Biological function summary	MRC2 serves several significant roles including its involvement in tissue remodeling cellular migration and extracellular matrix turnover. The protein forms part of a larger complex involving other matrix-interacting proteins allowing for efficient capture and presentation of ligands. MRC2 contributes to the regulation

	of collagen homeostasis through interaction with collagenase and facilitates uptake and processing of collagen.
Pathways	MRC2 takes part in cellular processes connected to collagen metabolism and tissue remodeling. It is particularly involved in pathways such as the receptor-mediated endocytosis pathway and collagen degradation pathway. Interactions between MRC2 and proteins like uPAR (urokinase receptor) highlight its role in plasminogen activation which is essential for extracellular matrix breakdown and cell migration.
Associated diseases and disorders	MRC2 displays connection to tissue fibrosis and cancer metastasis. In fibrosis dysregulation of MRC2 can lead to excessive collagen deposition which contributes to disease progression. In cancer abnormal MRC2 activity associates with enhanced tumor cell invasion and metastasis. Additionally MRC2's interaction with uPAR suggests its involvement in tumor progression further emphasizing its potential as a therapeutic target for disease modulation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

8 product images



Immunocytochemistry/ Immunofluorescence - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)


This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized MG 63 (human osteosarcoma fibroblast) cells labelling MRC2/ENDO180 with ab317323 at 1/100 (4.98 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green).

Confocal image showing cytoplasmic staining in MG 63 cell line (shown in green). The counterstain was observed in magenta. Nuclear DNA was labeled with DAPI (shown in blue).
Negative control: MCF7 (PMID: 17974964).
Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

This data was developed using ab317323, the same antibody clone in a different buffer formulation.


Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue labeling MRC2/ENDO180 with ab317323 at 1/500 (0.996 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on stromal cells of human breast carcinoma
(PMID 26316068).
The section was incubated with ab317323 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling MRC2/ENDO180 with ab317323 at 1/500 (0.996 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Positive staining in endothelial cells of human kidney
(PMID: 11156692).
The section was incubated with ab317323 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

MRC2/ENDO180 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human cerebrum using rabbit Anti-MRC2/ENDO180 antibody

This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Immunohistochemical analysis of paraffin-embedded Human cerebrum tissue labeling MRC2/ENDO180 with ab317323 at 1/500 (0.996 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Low expression tissue: No staining on human cerebrum
(PMID: 8702911).
The section was incubated with ab317323 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins

---



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling MRC2/ENDO180 with ab317323 at 1/500 (0.996 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Positive staining in macrophages of human tonsil
(PMID: 8702911).

The section was incubated with ab317323 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Flow Cytometry - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Flow cytometric analysis of MCF7 (human breast adenocarcinoma epithelial cell, Left) / MG-63(human osteosarcoma fibroblast, Right) cells labelling MRC2/ENDO180 with ab317323 at 1/500 dilution (0.1 ug)/Red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Negative control: MCF7
Gated on viable cell.



Western blot - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

MRC2/ENDO180 Western blot staining using rabbit Anti-MRC2/ENDO180 antibody

This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Low expression: MCF7, HepG2.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-MRC2/ENDO180 antibody [EPR29048-48] (ab317323) at 1/1000 dilution

Lane 1:
Saos-2 (human osteosarcoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
MG-63 (human osteosarcoma fibroblast) whole cell lysate at 20 µg

Lane 3:
MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 4:
HepG2 (human hepatocellar carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 5:

U-2 OS (human bone osteosarcoma epithelial cell) whole cell lysate at 20 µg


Lane 6:
A-172 (human brain glioblastoma cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 180 kDa, 36 kDa

Exposure time: 6s



Western blot - Anti-MRC2/ENDO180 antibody [EPR29048-48] - BSA and Azide free (ab317324)

MRC2/ENDO180 Western blot staining using rabbit Anti-MRC2/ENDO180 antibody

This data was developed using ab317323, the same antibody clone in a different buffer formulation.

Low expression tissue: human brain (PMID: 8702911).

The identity of the lower MW band at approximately 65 kDa (in lane 3) is unknown.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

Exposure time: Lane 1: 103 seconds; Lanes 2-3: 180 seconds.

All lanes:
Western blot - Anti-MRC2/ENDO180 antibody [EPR29048-48] (ab317323) at 1/1000 dilution

Lane 1:
Human liver cancer tissue lysate at 20 µg

Lane 2:
Human tonsil tissue lysate at 20 µg

Lane 3:
Human brain tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Observed band size: 180 kDa, 36 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.