abcam

ab317845

Anti-Dnmt1 antibody [RM1192]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Multiclonal
Immunogens	The exact immunogen used to generate this antibody is proprietary information.
Clone number	RM1192
Purification technique	Affinity purification Protein A
Specificity	Unsuitable for mouse IP and rat ICC.
Concentration	0.531 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/500

Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/5000

Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information DNMT1

Function	DNA methyltransferase that methylates CpG residues (PubMed:17200670, PubMed:18754681, PubMed:21745816, PubMed:26070743). Preferentially methylates hemimethylated DNA (PubMed:21745816, PubMed:26070743). Associates with DNA replication sites in S phase maintaining the methylation pattern in the newly synthesized strand, that is essential for epigenetic inheritance (PubMed:17200670, PubMed:21745816). Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication (PubMed:21745816). It is responsible for maintaining methylation patterns established in development (PubMed:21745816). DNA methylation is coordinated with methylation of histones (PubMed:16357870). Mediates transcriptional repression by direct binding to HDAC2 (PubMed:10888872). In association with DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9 (PubMed:18413740). Probably forms a corepressor complex required for activated KRAS-mediated promoter hypermethylation and transcriptional silencing of tumor suppressor genes (TSGs) or other tumor-related genes in colorectal cancer (CRC) cells (PubMed:24623306). Also required to maintain a transcriptionally repressive state of genes in undifferentiated embryonic stem cells (ESCs) (PubMed:24623306). Associates at promoter regions of tumor suppressor genes (TSGs) leading to their gene silencing (PubMed:24623306).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What are recombinant multiclonals?
Recombinant multiclonals are a mixture of recombinant antibodies co-expressed from a library of heavy and light chains. They offer several advantages including:

- The sensitivity of polyclonal antibodies by recognising multiple epitopes
- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

View our range of recombinant multiclonal antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Dnmt1 antibody [RM1192] (ab317845)


Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling Dnmt1 with ab317845 at 1/500 (1.062 ug/ml) dilution, followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880).

Positve staining on human tonsil. The section was incubated with ab317845 at 4°C overnight.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880).

Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Dnmt1 antibody [RM1192] (ab317845)


Immunohistochemical analysis of paraffin-embedded A Wild-type HAP1 (human chronic myelogenous leukemia near-haploid cell) cell pellet. B Dnmt1 knockout HAP1 cell pellet. tissue labeling Dnmt1 with ab317845 at 1/500 (1.062 ug/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on (A) wild-type HAP1 cell pellet, no staining on (B) Dnmt1 knockout HAP1 cell pellet. The section was incubated with ab317845 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins



Flow Cytometry (Intracellular) - Anti-Dnmt1 antibody [RM1192] (ab317845)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Parental HAP1 (HOWT01) (human chronic myelogenous leukemia near-haploid cell) (Left) / DNMT1 knockout HAP1(HO010163) (Right) cells labelling Dnmt1 with ab317845 at 1/5000 dilution (0.1ug) (Magenta) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.




Immunocytochemistry/ Immunofluorescence - Anti-Dnmt1 antibody [RM1192] (ab317845)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized parental HAP1(HOWT01) (human chronic myelogenous leukemia near-haploid cell) cells labelling Dnmt1 with ab317845 at 1/500 (1.062 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green).

Confocal image showing nuclear staining in wildtype HAP1 cell line (shown in green) and no staining in DNMT1 KO HAP1 cell line. The counterstain was observed in magenta. Nuclear DNA was labeled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Immunoprecipitation - Anti-Dnmt1 antibody [RM1192] (ab317845)

Dnmt1 was immunoprecipitated from 0.35 mg HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate with ab317845 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab317845 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate
Lane 2: ab317845 IP in HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab317845 in HeLa whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST

The bands beneath the target band are likely to be degraded target fragments.


All lanes:
Immunoprecipitation - Anti-Dnmt1 antibody [RM1192] (ab317845) at 1/30 dilution

All lanes:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 8s



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Dnmt1 antibody [RM1192] (ab317845)


Immunohistochemical analysis of paraffin-embedded Rat testis tissue labeling Dnmt1 with ab317845 at 1/100 (5.31 ug/ml) dilution, followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880).

Positive staining on rat testis. The section was incubated with ab317845 at 4°C overnight.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880).

Heat mediated antigen retrieval was performed using ab93678 (citrate buffer, pH 6.0)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Dnmt1 antibody [RM1192] (ab317845)


Immunohistochemical analysis of paraffin-embedded Mouse testis tissue labeling Dnmt1 with ab317845 at 1/100 (5.31 ug/ml) dilution, followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880).

Positive staining on mouse testis. The section was incubated with ab317845 at 4°C overnight.

Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880).

Heat mediated antigen retrieval was performed using ab93678 (citrate buffer, pH 6.0)




Immunocytochemistry/ Immunofluorescence - Anti-Dnmt1 antibody [RM1192] (ab317845)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Neuro-2a (mouse neuroblastoma neuroblast) cells labelling Dnmt1 with ab317845 at 1/500 (1.062 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green).

Confocal image showing nuclear staining in neuro-2a cell line (shown in green). The counterstain was observed in magenta. Nuclear DNA was labeled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 31426844).

The lysates were freshly made and used for Western Blotting immediately to minimize protein degradation.

Exposure time: Lane 1-2: 10 seconds Lane 2: 70 seconds

All lanes:

Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845) at 1/1000 dilution

Lane 1:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg


Lane 2:
293T (human embryonic kidney epithelial cell) whole cell lysate at 20 µg

Lane 3:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 183 kDa, 184 kDa



Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 31426844).

The lysates were suggested freshly made and used for Western Blotting immediately to minimize protein degradation.

Exposure time: Lane 1:180 seconds Lane 2: 70 seconds

All lanes:
Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845) at 1/1000 dilution


Lane 1:
Frozen HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Fresh HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 183 kDa, 184 kDa



Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845)

Blocking and diluting buffer and concentration: Intercept® (TBS) Blocking Buffer diluted with an equal volume of TBS.

Lysates at 20 µg per lane.

The samples were run on a Bis-Tris gel under reducing conditions.

Western blot: Anti-Dnmt1 antibody (ab317845) staining at 1/1000 dilution and Anti-Dnmt1 antibody (ab19905) staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in magenta.

In Western blot, ab317845 was shown to bind specifically to Dnmt1. Target of interest was observed at 183-184 kDa in wild-type HAP1 cell lysates (lane 1) with no signal observed at this size in Dnmt1 knockout cell line (lane 2). To generate this image, samples were first run on an SDS-PAGE gel then transferred onto an immobilon-FL PVDF membrane. Membranes were blocked in a fluorescent western blot (TBS-based) blocking solution before incubation with primary antibodies overnight at 4 °C. Blots were washed in TBS-T, incubated with secondary antibodies Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution for 1 h at room temperature, washed again then imaged.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 31426844).
All lanes:
Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845) at 1/1000 dilution

Lane 1:
Wild-type HAP1 (human chronic myelogenous leukemia near-haploid cell) whole cell lysate at 20 µg

Lane 2:
Dnmt1 knockout HAP1 whole cell lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG H&L (800CW) and Goat Anti-Mouse IgG H&L (680RD) at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 183 kDa, 184 kDa, 36 kDa

Exposure time: 180s



Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 31426844).

The lysates were freshly made and used for Western Blotting immediately to minimize protein degradation.

All lanes:
Western blot - Anti-Dnmt1 antibody [RM1192] (ab317845) at 1/1000 dilution

Lane 1:
Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Lane 2:
SH-SY5Y (human neuroblastoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 183 kDa, 184 kDa

Exposure time: 59s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.