abcam

ab318166

Alexa Fluor® 488 Anti-Tissue Factor antibody [EPR22548-232]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR22548-232
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

IHC-P

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information F3

Function	Initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The [TF:VIIa] complex activates factors IX or X by specific limited proteolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Tissue Factor also known as coagulation factor III or CD142 is an important player in the blood coagulation process. This protein with a molecular mass of approximately 47 kDa is a transmembrane glycoprotein. It is expressed widely in various tissues including the lungs heart and brain and plays an important role in initiating the coagulation cascade when exposed to blood. In endothelial cells tissue factor is generally not present unless these cells are in a state of disturbance supporting its role as a primary defense mechanism against vascular injury.

Biological function summary	Tissue factor acts as a cofactor for factor VIIa forming a complex that rapidly catalyzes the activation of factor X to factor Xa. This catalytic action ultimately leads to fibrin formation which is essential for clotting. Tissue Factor (TF) is part of the extrinsic pathway of the coagulation cascade and is fundamental in the generation of a hemostatic response to tissue injury. By regulating thrombin generation through its interaction with other coagulation factors tissue factor plays an important role in maintaining hemostatic balance.
Pathways	Tissue factor is integral in the extrinsic pathway of coagulation which works closely with the intrinsic pathway to ensure effective hemostasis. It complexes with factor VIIa and triggers downstream effects involving proteins like factor X and thrombin. Tissue factor forms a bridge between cellular injury responses and the blood coagulation system linking these two significant physiological processes. The interaction with these proteins highlights its central role in hemostatic pathways and highlights its significance in the coagulation network.
Associated diseases and disorders	Tissue factor is associated with conditions like thrombosis and cardiovascular diseases where its expression can become dysregulated. Abnormal activation of tissue factor contributes directly to the pathological clot formation seen in thrombosis. It is also implicated in cancer-related coagulopathy where it can facilitate tumor-associated thrombosis by interacting with cancer cell surfaces. Additionally in such diseases the interplay between tissue factor and thrombin becomes important as it affects disease progression and the severity of clot-related complications.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:

1 product image



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-Tissue Factor antibody [EPR22548-232] (ab318166)

Tissue Factor Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Tissue Factor antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized A431 (human epidermoid carcinoma epithelial cell) cells labelling Tissue Factor with ab318166 at 1/50 (10.0 ug/ml) dilution (Green).

Confocal image showing membranous staining in A431 cell line (shown in green). The counterstain was observed in magenta. Nuclear DNA was labeled with DAPI (shown in blue). Low expression: MCF7 (PMID: 24137414, 28938620). Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.