abcam

ab319039

Anti-CD200R antibody [EPR28088-189]

Anti-CD200R antibody [EPR28088-189] is a Rabbit Monoclonal antibody that is used in CD200R Flow Cytometry, Western Blot. Suitable for Human samples.

This gene encodes a receptor for the OX-2 membrane glycoprotein. Both the receptor and substrate are cell surface glycoproteins containing two immunoglobulin-like domains. This receptor is restricted to the surfaces of myeloid lineage cells and the receptor-substrate interaction may function as a myeloid downregulatory signal.

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR28088-189
Purification technique	Affinity purification Protein A
Concentration	0.505 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Human, Mouse, Rat

Dilution info	-
Notes	-

Target data

See full target information CD200R1

Function	Inhibitory receptor for the CD200/OX2 cell surface glycoprotein. Limits inflammation by inhibiting the expression of pro-inflammatory molecules including TNF, interferons, and inducible nitric oxide synthase (iNOS) in response to selected stimuli. Also binds to HHV-8 K14 viral CD200 homolog with identical affinity and kinetics as the host CD200.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD200R also known as CD200 receptor is a protein expressed mainly on myeloid cells including macrophages dendritic cells and certain T cells. The protein has a molecular mass of approximately 62 kDa. This receptor is a type I membrane glycoprotein that plays an important role in immune system regulation. It binds to its ligand CD200 protein found on various cell types such as neurons epithelial cells and some immune cells to mediate suppressive effects on the immune response.
Biological function summary	CD200R plays a significant role in controlling the immune response by acting as an inhibitory receptor. When activated it sends signals that dampen immune cell activity which can prevent overactive immune responses and promote tissue homeostasis. This receptor is not part of a large receptor complex but interacts directly with its ligand CD200 to initiate signaling cascades that modulate immune functions. Its primary role lies in preventing unnecessary inflammation and autoimmunity through negative regulation of macrophage activation.
Pathways	CD200R is involved in the regulatory pathways of the immune system. It participates in pathways that manage immune tolerance and its interactions impact pathways related to inflammation control and tissue repair. The CD200R signaling cascades intersect with pathways involving other immune-modulatory proteins such as CD200R1 and CD200R3 which contribute to cellular communication in the immune environment. These pathways play critical roles in downregulating inflammatory responses and maintaining immune balance.
Associated diseases and disorders	CD200R holds relevance for conditions involving immune dysregulation. It is connected to autoimmune disorders and atopic dermatitis. For example in atopic dermatitis altered expression or function of CD200R may lead to improper immune responses contributing to inflammation. Furthermore CD200R agonists which are molecules that mimic the function of CD200 are an area of interest for developing therapeutic strategies to modulate immune responses. Related proteins such as CD200 and its other receptor variants play active roles in these disorders highlighting the importance of their balanced function to maintain immune homeostasis.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

5 product images




Flow Cytometry - Anti-CD200R antibody [EPR28088-189] (ab319039)

Flow cytometric analysis of No-GFP-CD16.NK-92 (CD16 ectopically expressed natural killer(NK-92®) cell with no GFP tag) (Right) / A549 (human lung carcinoma epithelial cell) (Left) cells labelling CD200R with ab319039 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Low expression: A549.



Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The identity of the bands lower than 50 kDa are unknown.

All lanes:
Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039) at 1/1000 dilution

Lane 1:
HEL (human erythroleukemia erythroblast) whole cell lysate at 20 µg


Lane 2:
HuT-78 (human sezary syndrome cutaneous t lymphocyte) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 60-90 kDa

Exposure time: 180s



Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Low expression: A549, NCI-H1299 (PMID: 32395395).

The identity of the bands lower than 50 kDa are unknown.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039) at 1/1000 dilution

Lane 1:
A549 (human lung carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
No-GFP-CD16.NK-92 (natural killer (nk-92®) cell lacki gfp-cd16) whole cell lysate at 20 µg


Lane 3:
NCI-H1299 (human lung carcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 60-90 kDa, 36 kDa

Exposure time: 10s



Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The identity of the lower MW band at approximately 48 kDa is unknown.

The high-sensitivity ECL substrate allows for the detection of proteins in the mid-femtogram range.

All lanes:
Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039) at 1/1000 dilution

All lanes:
Human placenta tissue lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Developed using the ECL technique.

Observed band size: 60-90 kDa

Exposure time: 136s



Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Low expression: skeletal muscle (PMID: 12853143).

The identity of the lower MW band at approximately 36 kDa is unknown.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lanes:
Western blot - Anti-CD200R antibody [EPR28088-189] (ab319039) at 1/1000 dilution

Lane 1:
Human spleen tissue lysate at 20 µg

Lane 2:
Human skeletal muscle tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Observed band size: 60-75 kDa, 36 kDa

Exposure time: 136s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.