

Anti-AMPK alpha 1 antibody [Y365]

Anti-AMPK alpha 1 antibody [Y365] (ab32047) is a rabbit monoclonal antibody detecting AMPK alpha 1 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 190 publications
- Trusted since 2006

Lab Essentials

Recombinant

RabMAb

KO Validated

20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	Y365
Purification technique	Affinity purification Protein A
Specificity	This antibody is specific for human AMPK alpha 1. This antibody shows low sensitivity in mouse and rat samples.
Concentration	0.76 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/40 - 1/50
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	African green monkey
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000 - 1/5000
Notes	This antibody shows low sensitivity in mouse and rat samples.

Species	Rat
Dilution info	1/1000 - 1/5000
Notes	This antibody shows low sensitivity in mouse and rat samples.

Species	Human
Dilution info	1/1000 - 1/5000
Notes	This antibody shows low sensitivity in mouse and rat samples.

Predicted

Species	African green monkey
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	African green monkey
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/150
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
---------	------------

Dilution info	Use at an assay dependent concentration.
----------------------	--

Notes	-
--------------	---

Predicted

Species	African green monkey
----------------	----------------------

Dilution info	-
----------------------	---

Notes	-
--------------	---

IHC-P

Tested

Species	Human
----------------	-------

Dilution info	1/100 - 1/250
----------------------	---------------

Notes	Mouse and rat species are recommended by WB application. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
--------------	--

Expected

Species	Mouse, Rat
----------------	------------

Dilution info	Use at an assay dependent concentration.
----------------------	--

Notes	-
--------------	---

Predicted

Species	African green monkey
----------------	----------------------

Dilution info	-
----------------------	---

Notes	-
--------------	---

Target data

[See full target information PRKAA1](#) 

Function	Catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism (PubMed:17307971, PubMed:17712357, PubMed:24563466, PubMed:31492851,
-----------------	---

PubMed:37821951, PubMed:40233740). In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation (PubMed:17307971, PubMed:17712357). AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators (PubMed:17307971, PubMed:17712357). Regulates lipid synthesis by phosphorylating and inactivating lipid metabolic enzymes such as ACACA, ACACB, GYS1, HMGCR and LIPE; regulates fatty acid and cholesterol synthesis by phosphorylating acetyl-CoA carboxylase (ACACA and ACACB) and hormone-sensitive lipase (LIPE) enzymes, respectively (By similarity). Promotes lipolysis of lipid droplets by mediating phosphorylation of isoform 1 of CHKA (CHKalpha2) (PubMed:34077757). Regulates insulin-signaling and glycolysis by phosphorylating IRS1, PFKFB2 and PFKFB3 (By similarity). AMPK stimulates glucose uptake in muscle by increasing the translocation of the glucose transporter SLC2A4/GLUT4 to the plasma membrane, possibly by mediating phosphorylation of TBC1D4/AS160 (By similarity). Regulates transcription and chromatin structure by phosphorylating transcription regulators involved in energy metabolism such as CRTC2/TORC2, FOXO3, histone H2B, HDAC5, MEF2C, MLXIPL/ChREBP, EP300, HNF4A, p53/TP53, SREBF1, SREBF2 and PPARGC1A (PubMed:11518699, PubMed:11554766, PubMed:15866171, PubMed:17711846, PubMed:18184930). Acts as a key regulator of glucose homeostasis in liver by phosphorylating CRTC2/TORC2, leading to CRTC2/TORC2 sequestration in the cytoplasm (By similarity). In response to stress, phosphorylates 'Ser-36' of histone H2B (H2BS36ph), leading to promote transcription (By similarity). Acts as a key regulator of cell growth and proliferation by phosphorylating FNIP1, TSC2, RPTOR, WDR24 and ATG1/ULK1: in response to nutrient limitation, negatively regulates the mTORC1 complex by phosphorylating RPTOR component of the mTORC1 complex and by phosphorylating and activating TSC2 (PubMed:14651849, PubMed:18439900, PubMed:20160076, PubMed:21205641). Also phosphorylates and inhibits GATOR2 subunit WDR24 in response to nutrient limitation, leading to suppress glucose-mediated mTORC1 activation (PubMed:36732624). In response to energetic stress, phosphorylates FNIP1, inactivating the non-canonical mTORC1 signaling, thereby promoting nuclear translocation of TFEB and TFE3, and inducing transcription of lysosomal or autophagy genes (PubMed:37079666). In response to nutrient limitation, promotes autophagy by phosphorylating and activating ATG1/ULK1 (PubMed:21205641). In that process, it also activates WDR45/WIPI4 (PubMed:28561066). Phosphorylates CASP6, thereby preventing its autoprocessing and subsequent activation (PubMed:32029622). In response to nutrient limitation, phosphorylates transcription factor FOXO3 promoting FOXO3 mitochondrial import (By similarity). Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin (PubMed:17486097). AMPK also acts as a regulator of circadian rhythm by mediating phosphorylation of CRY1, leading to destabilize it (By similarity). May regulate the Wnt signaling pathway by phosphorylating CTNNB1, leading to stabilize it (By similarity). Also has tau-protein kinase activity: in response to amyloid beta A4 protein (APP) exposure, activated by CAMKK2, leading to phosphorylation of MAPT/TAU; however the relevance of such data remains unclear in vivo (By similarity). Also phosphorylates CFTR, EEF2K, KLC1, NOS3 and SLC12A1 (PubMed:12519745, PubMed:20074060). Regulates hepatic lipogenesis. Activated via SIRT3, represses sterol regulatory element-binding protein (SREBP) transcriptional activities and ATP-consuming lipogenesis to restore cellular energy balance. Upon stress, regulates mitochondrial fragmentation through phosphorylation of MTFR1L (PubMed:36367943). Phosphorylates ALDH7A1 in response to cellular stress, such

as hypoxia or ferroptotic stress, promoting ALDH7A1 recruitment to membranes (PubMed:31492851, PubMed:40233740).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Anti-AMPK alpha 1 antibody [Y365] (ab32047) was developed by Abcam using patented rabbit monoclonal antibody technology and is validated for use in Flow Cyt (Intra), ICC/IF, IHC-P, IP, WB in human, mouse, rat samples.

Anti-AMPK alpha 1 antibody [Y365] (ab32047) has been cited over 192 times in peer reviewed journals and is trusted by the scientific community.

Abcam's high quality manufacturing and validation processes ensure Anti-AMPK alpha 1 antibody [Y365] (ab32047) has high sensitivity and specificity alongside high lot-to-lot consistency and reproducibility.

The specificity of Anti-AMPK alpha 1 antibody [Y365] (ab32047) has been confirmed by testing in knockout samples.

Anti-AMPK alpha 1 antibody [Y365] (ab32047) has 5 independent reviews from customers.

Anti-AMPK alpha 1 antibody [Y365] (ab32047) specifically detects AMPK alpha 1 (UniProt ID: Q13131; Molecular weight: 64kDa) and is sold in a convenient trial size to enable initial testing (20 µL) and larger sizes for subsequent scaling up experiments (100 µL and 1 mL).

Conjugation-ready, carrier free format available for antibody clone Y365 - ab210714.

Antibody clone Y365 is also available pre-conjugated to a variety of labels for your convenience - PE, APC, HRP, Alkaline Phosphatase, Alexa Fluor® 488, Alexa Fluor® 647, Alexa Fluor® 594, Alexa Fluor® 555 (ab303053, ab303054, ab303055, ab308710, ab309679, ab310041, ab310429, ab311955).

AMPK alpha 1/AMPKα1 (encoded by the PRKAA1 gene) is a key regulator of cellular energy metabolism and metabolic regulation. It plays a crucial role in maintaining energy homeostasis by activating pathways that generate ATP and inhibiting those that consume it. Abnormal AMPK alpha 1 activity is linked to diseases such as cancer and diabetes,

highlighting its disease relevance. Our AMPKa1 antibody is a valuable tool for studying these pathways and understanding the molecular mechanisms involved in these conditions.

Patented technology

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

AMPK alpha 1 Western blot staining using rabbit Anti-AMPK alpha 1 antibody

False colour image of Western blot: Anti-AMPK alpha 1 antibody [Y365] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab32047 was shown to bind specifically to AMPK alpha 1. A band was observed at 64 kDa in wild-type RAW 264.7 cell lysates with no signal observed at this size in PRKAA1 knockout cell line [ab280055](#) (knockout cell lysate [ab280114](#)). To generate this image, wild-type and PRKAA1 knockout RAW 264.7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.

All lanes:

Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/1000 dilution

Lane 1:

Wild-type RAW 264.7 cell lysate at 20 µg

Lane 2:

Western blot - Mouse PRKAA1 knockout RAW 264.7 cell lysate ([ab280114](#)) at 20 µg

Lane 3:

Mouse Liver cell lysate at 20 µg


Lane 4:

Neuro2A cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 64 kDa


Observed band size: 64 kDa



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)


AMPK alpha 1 Flow Cytometry (Intracellular) staining using rabbit Anti-AMPK alpha 1 antibody

Overlay histogram showing HeLa cells stained with unpurified ab32047 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (unpurified ab32047, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervical carcinoma tissue labelling AMPK alpha 1 with unpurified ab32047 at a dilution of 1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

AMPK alpha 1 Flow Cytometry (Intracellular) staining using rabbit Anti-AMPK alpha 1 antibody


Intracellular Flow Cytometry analysis of HeLa cells labelling AMPK alpha 1 with purified ab32047 at 1/150 (red). Cells were fixed with 2% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

AMPK alpha 1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-AMPK alpha 1 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling AMPK alpha 1 with purified ab32047 at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9 before commencing with IHC staining protocol. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.




Immunocytochemistry/ Immunofluorescence - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

AMPK alpha 1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-AMPK alpha 1 antibody

Immunocytochemistry/Immunofluorescence analysis of MCF7 cells labelling AMPK alpha 1 with purified ab32047 at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. [ab150077](#), an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. [ab7291](#), a mouse anti-tubulin (1/1000) and [ab150120](#), an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/250) and secondary antibody, [ab150120](#), an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: [ab7291](#) (1/1000) and secondary antibody, [ab150077](#), an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).



Immunoprecipitation - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

ab32047 (purified) at 1/40 immunoprecipitating AMPK alpha 1 in HeLa whole cell lysate.

Lane 1 (input): HeLa whole cell lysate (10µg)

Lane 2 (+): ab32047 + HeLa whole cell lysate (10µg).

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab32047 in HeLa whole cell lysate.

For western blotting, [ab131366](#) VeriBlot for IP Detection Reagent (HRP) was used for detection (1/1500).

Blocking buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM /TBST.

All lanes:

Immunoprecipitation - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Predicted band size: 64 kDa

Observed band size: 63 kDa



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

This antibody shows low sensitivity in mouse and rat samples.

All lanes:

Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/20000 dilution

Lane 1:

MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:

HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:

HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 4:

K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate at 20 µg

Lane 5:

C6 (Rat glial tumor glial cell) whole cell lysate at 20 µg

Lane 6:

Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Lane 7:

NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate at 20 µg

Secondary


All lanes:

Western blot - Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 64 kDa

Observed band size: 63 kDa

Exposure time: 180s



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

AMPK alpha 1 Western blot staining using rabbit Anti-AMPK alpha 1 antibody

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: AMPK alpha knockout HAP1 cell lysate (20 µg)

Lane 3: MCF7 cell lysate (20 µg)

Lane 4: HeLa cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab32047 observed at 63 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab32047 was shown to specifically react with AMPK alpha in wild-type HAP1 cells. No band was observed when AMPK alpha knockout samples were examined. Wild-type and AMPK alpha knockout samples were subjected to SDS-PAGE. ab32047 and [ab8245](#) (loading control to GAPDH) were diluted 1/5000 and 1/2000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10,000 dilution for 1hr at room temperature before imaging.

All lanes:

Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Predicted band size: 64 kDa



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Blocking/Diluting buffer and concentration: 5% NFDm/TBST.

This antibody shows low sensitivity in mouse samples.

All lanes:

Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/500 dilution

Lane 1:

HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate at 20 µg

Lane 2:

NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 3:

Mouse liver tissue lysate at 20 µg

Lane 4:

Mouse brain tissue lysate at 20 µg

Lane 5:

Mouse kidney tissue lysate at 20 µg

Lane 6:

Mouse retina tissue lysate at 20 µg

Lane 7:

Mouse skeletal muscle tissue lysate at 20 µg

Secondary


All lanes:

Western blot - Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 64 kDa

Observed band size: 40 kDa, 63 kDa

Exposure time: 3min



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Blocking and diluting buffer and concentration: 5% NFDM /TBST.

[ab181602](#) was used as a GAPDH loading control at 1/1000000 dilution.

This blot was developed using a higher sensitivity ECL substrate.

Compared with ab32047, [ab271188](#) has higher sensitivity, we recommend [ab271188](#) as an alternative for testing AMPK alpha 1 in mouse and rat samples in western blot.

Lanes 1 - 8:

Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/1000 dilution

Lanes 1 - 8:

Western blot - Anti-AMPK alpha 1 antibody [EPR24413-70] ([ab271188](#)) at 1/1000 dilution

Lane 1:

NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 2:

Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Lane 3:

C6 (rat glial tumor glial cell) whole cell lysate at 20 µg

Lane 4:

Mouse brain tissue lysate at 20 µg

Lane 5:

Mouse heart tissue lysate at 20 µg

Lane 6:

Mouse skeletal muscle tissue lysate at 20 µg

Lane 7:

Rat brain tissue lysate at 20 µg

Lane 8:

Rat heart tissue lysate at 20 µg

Secondary


All lanes:

Western blot - Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Developed using the ECL technique.

Observed band size: 64 kDa

Exposure time: 180s



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

AMPK alpha 1 Western blot staining using rabbit Anti-AMPK alpha 1 antibody

Blocking and dilution buffer: 5% NFDM/TBST.

This antibody shows low sensitivity in mouse and rat samples.

All lanes:

Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/2000 dilution

Lane 1:

C6 cell lysate at 20 µg

Lane 2:

NIH/3T3 cell lysate at 20 µg

Secondary

All lanes:

Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

Predicted band size: 64 kDa

Observed band size: 63 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.