abcam

ab32049

Anti-p53 antibody [Y5]

Anti-p53 antibody [Y5] (ab32049) is a rabbit monoclonal antibody detecting p53 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 50 publications
- Trusted since 2006

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	Y5
Purification technique	Affinity purification Protein A
Specificity	This antibody clone recognises both wild-type and mutant forms of p53 in human samples. It is not designed to recognise any specific p53 mutation. We have confirmed this experimentally and have been able to detect p53 in different cell lines using various applications and treatments. Important note: p53 expression levels vary greatly between cell lines. It has been reported that p53 mutations render the protein more stable, hence mutated cell lines often express higher levels of the p53 protein compared to wild-type cell lines. For low expressing wild type cell lines, p53 expression can be increased with cell treatments such as camptothecin or irinotecan.
Dissociation constant	2.02 x 10^{-10} M
Concentration	0.106 mg/mL The concentration of this product may be batch-dependent

Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

IP

Tested

Species	Human
Dilution info	1/20 for 2 µg
Notes	-

WB

Tested

Species	Human
Dilution info	1/500 - 1/5000
Notes	This antibody recognises both wild-type and mutant forms of human p53 and is not specific to any particular mutation. Expression levels of p53 can vary significantly between cell lines, with mutant forms often showing higher stability and abundance. In low-expressing wild-type cell lines, p53 levels may be increased using treatments such as camptothecin or irinotecan.

ICC/IF

Tested

Species	Human
Dilution info	1/100

Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information TP53

Function

Multifunctional transcription factor that induces cell cycle arrest, DNA repair or apoptosis upon binding to its target DNA sequence (PubMed:11025664, PubMed:12524540, PubMed:12810724, PubMed:15186775, PubMed:15340061, PubMed:17317671, PubMed:17349958, PubMed:19556538, PubMed:20673990, PubMed:20959462, PubMed:22726440, PubMed:24051492, PubMed:24652652, PubMed:35618207, PubMed:36634798, PubMed:38653238, PubMed:9840937). Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type (PubMed:11025664, PubMed:12524540, PubMed:12810724, PubMed:15186775, PubMed:15340061, PubMed:17189187, PubMed:17317671, PubMed:17349958, PubMed:19556538, PubMed:20673990, PubMed:20959462, PubMed:22726440, PubMed:24051492, PubMed:24652652, PubMed:38653238, PubMed:9840937). Negatively regulates cell division by controlling expression of a set of genes required for this process (PubMed:11025664, PubMed:12524540, PubMed:12810724, PubMed:15186775, PubMed:15340061, PubMed:17317671, PubMed:17349958, PubMed:19556538, PubMed:20673990, PubMed:20959462, PubMed:22726440, PubMed:24051492, PubMed:24652652, PubMed:9840937). One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression (PubMed:12524540, PubMed:17189187). Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed:12524540). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed:12524540). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by

repressing CLOCK-BMAL1-mediated transcriptional activation of PER2 (PubMed:24051492).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-p53 antibody [Y5] (ab32049) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human samples.

What is the molecular weight of p53?
Anti-p53 [Y5] (ab32049) specifically detects a band for p53 (UniProt: P04637) at a molecular weight of 44kDa.

Trusted by the scientific community
Anti-p53 [Y5] (ab32049) was first used in a scientific publication in 2006 and has been cited over 50 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-p53 antibody [Y5] (ab32049) has been confirmed by Western blot testing in TP53 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [Y5] also available for your convenience: ab32049, HRP - ab190335, Carrier free - ab219731, Alexa Fluor® 488 - ab224920, APC - ab310851, PE - ab310921, Alexa Fluor® 647 - ab311085, Alexa Fluor® 594 - ab311682, Alexa Fluor® 568 - ab312957, Alexa Fluor® 555 - ab313166, Alexa Fluor® 750 - ab321560

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Species reactivity
Rat: We have preliminary internal testing data to indicate this antibody may not react with this species.
Please contact us for more information.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The protein p53 also known as TP53 or tumor protein p53 has a molecular weight of approximately 53 kDa. It acts as a transcription factor and plays a major role in cell cycle regulation apoptosis and maintaining genomic stability. This protein mainly expresses in the nucleus of cells and acts as a critical regulator of cellular responses to stress signals including DNA damage. Scientists commonly use p53 antibodies in various assays like western blot and p53 immunofluorescence to detect and study its expression and functional status in cells.
Biological function summary	P53 functions to control cell division and apoptosis serving as a guardian of the genome by preventing mutation accumulation. It does not form part of a larger complex under normal conditions but interacts with various other molecules to execute its functions. p53 can activate or suppress the transcription of numerous genes involved in cell cycle arrest DNA repair and programmed cell death allowing it to halt the progression of damaged cells and trigger repair mechanisms or eliminate those that cannot be repaired.
Pathways	P53 acts within several key biological pathways such as the p53 signaling pathway and the intrinsic apoptotic pathway. Its activity involves interaction with proteins like MDM2 which regulates p53 through ubiquitin-mediated degradation and ATM kinase which phosphorylates p53 in response to DNA damage. These interactions ensure appropriate cellular responses during stress and are vital for maintaining homeostasis.
Associated diseases and disorders	P53 mutation or inactivation is often associated with the development of cancer given its role in controlling cell division and preventing tumor formation. Specifically its dysfunction has been linked to cancers such as breast cancer and lung cancer. Additionally p53 can interact with other mutant proteins like Ras compounding mutations that contribute to tumor progression and aggressive cancer phenotypes. Understanding these interactions and the status of p53 can be important in developing targeted cancer therapies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


23 product images



Flow Cytometry (Intracellular) - Anti-p53 antibody [Y5] (ab32049)

P53 Flow Cytometry (Intracellular) staining using rabbit Anti-p53 antibody


Flow cytometry overlay histogram showing p53 in wild-type HAP1 (green line) and TP53 knockout HAP1 (red line) cells stained with ab32049. The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab32049) (1x 106 cells in 100µl at 0.2 µg/ml (1/440)) for 30min at 22°C. The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed (ab150081) was incubated at 1/4000 for 30min at 22°C. Isotype control antibody Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) was used at the same concentration and conditions as the primary antibody (wild-type HAP1 - black line, TP53 knockout HAP1 - grey line). Unlabelled sample was also used as a control (this line is not shown for the purpose of simplicity). Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. This antibody gave a positive signal in HAP1 Fixed with 80% methanol (5 min) / permeabilised with 0.1% PBS-Triton X-100 for 15 min under the same conditions.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody


Immunohistochemistry (Paraffin-embedded sections) using ab32049 at a dilution of 1/50 and human skin cancer
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody

Immunohistochemical analysis of paraffin embedded normal Human uterus tissue (negative control) labeling p53 with ab32049.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody


ab32049 showing positive staining in Breast carcinoma tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-p53 antibody [Y5] (ab32049)

ab32049 staining wild-type p53 in MCF7 cells (a low expressing cell line). The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab32049 at 0.2µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and ab150120, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue). Also suitable in cells fixed with 4% paraformaldehyde (10 min). Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Immunocytochemistry/ Immunofluorescence - Anti-p53 antibody [Y5] (ab32049)


ab32049 staining wild-type p53 in Hek293 cells (a high expressing cell line). The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab32049 at 0.2µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and ab150120, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue). Also suitable in cells fixed with 4% paraformaldehyde (10 min). Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Flow Cytometry (Intracellular) - Anti-p53 antibody [Y5] (ab32049)

P53 Flow Cytometry (Intracellular) staining using rabbit Anti-p53 antibody

Flow cytometry overlay histogram showing wild-type p53 in Hek-293 positive cells (left) and MCF7 negative cells (right) stained with ab32049 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab32049) (1x 106 cells in 100µl at 0.008µg/ml (1/13250)) for 30min at 22°C. The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed (ab150081) was incubated at 1/4000 for 30min at 22°C. Isotype control antibody (black line) Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) was used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control. Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. This antibody gave a positive signal in Hek-293 fixed with 80% methanol (5 min) / permeabilised with 0.1% PBS-Triton X-100 for 15 min under the same conditions.




Immunocytochemistry/ Immunofluorescence - Anti-p53 antibody [Y5] (ab32049)

ab32049 staining p53 in wild-type Hap1 cells (top panel) and p53 knockout Hap1 cells (bottom panel). The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at °C with ab32049 at 0.2µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and ab150120, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Also suitable in cells fixed with 100% methanol (5 min).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of wild-type HAP1 cells using rabbit Anti-p53 antibody

ab32049 was shown to react with TP53 in wild-type HAP1 cells in immunohistochemistry with loss of signal observed in a TP53 knockout cell line. HAP1 WT and TP53 KO cells were fixed and paraffin embedded. Sections were mounted onto slides and stained with ab32049 and the corresponding multimer HRP secondary. Chromogen image acquisition was performed, and representative images of each section are shown. Scale bar = 50 µm.

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody

ab32049 showing positive staining in Gastric adenocarcinoma tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody

ab32049 showing positive staining in Glioma tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody

ab32049 showing positive staining in Urinary bladder carcinoma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53 antibody [Y5] (ab32049)

P53 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p53 antibody


Immunohistochemical analysis of paraffin embedded normal Human breast tissue (negative control) labeling p53 with ab32049.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-p53 antibody [Y5] (ab32049)

P53 Flow Cytometry (Intracellular) staining using rabbit Anti-p53 antibody

Intracellular Flow Cytometry analysis of A431 (human epidermoid carcinoma) cells labeling p53 with unpurified ab32049 at 1/20 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) (1/2000) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control cells without incubation with primary antibody and secondary antibody (Blue) was used as the unlabeled control.




Immunocytochemistry/ Immunofluorescence - Anti-p53 antibody [Y5] (ab32049)

P53 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-p53 antibody

ab32049 staining p53 in A431 cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab32049 at 0.2µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and ab150120, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Immunoprecipitation - Anti-p53 antibody [Y5] (ab32049)

p53 was immunoprecipitated from 0.35 mg A431 (Human epidermoid carcinoma epithelial cell) whole cell lysate 10 µg with ab32049 at 1/100 dilution (2µg) . VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.

Lane 1: A431 (Human epidermoid carcinoma epithelial cell) whole cell lysate 10 µg

Lane 2: ab32049 IP in A431 whole cell lysate


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab32049 in A431 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-p53 antibody [Y5] (ab32049)

Predicted band size: 43 kDa

Observed band size: 44 kDa



Western blot - Anti-p53 antibody [Y5] (ab32049)

P53 Western blot staining using rabbit Anti-p53 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

In Western blot, Anti-p53 antibody [Y5] (ab32049) staining at 1/1000 dilution.

ab32049 recognises both wild-type and mutant forms of human p53 and is not specific to any particular mutation. Expression levels of p53 can vary significantly between cell lines, with mutant forms often showing higher stability and abundance. In low-expressing wild-type cell lines, p53 levels may be increased using treatments such as camptothecin or irinotecan.

All lanes:
Western blot - Anti-TP53 (phospho S376+S377+S392) antibody [RM1161] (ab322465) at 1/1000 dilution

Lane 1:
293T (human embryonic kidney epithelial cell) whole cell lysate (untreated membrane) at 20 µg


Lane 2:
293T whole cell lysate (alkaline phosphatase treated membrane) at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 53 kDa, 35-48 kDa, 36 kDa

Exposure time: 10s



Western blot - Anti-p53 antibody [Y5] (ab32049)

P53 Western blot staining using rabbit Anti-p53 antibody

Blocking/Diluting Buffer and concentration: 5% NFDM/TBST
Lane 1-4: Wildtype p53 cell lines

Lane 5-8: Mutant p53 cell lines

This antibody recognises both wild-type and mutant forms of human p53 and is not specific to any particular mutation. Expression levels of p53 can vary significantly between cell lines, with mutant forms often showing higher stability and abundance. In low-expressing wild-type cell lines, p53 levels may be increased using treatments such as camptothecin or irinotecan.

All lanes:
Western blot - Anti-p53 antibody [Y5] (ab32049) at 1/500 dilution

Lane 1:
HepG2 (Human hepatocellular carcinoma epithelial cell), Whole cell lysate at 20 µg

Lane 2:
A549 (Human lung carcinoma epithelial cell), Whole cell lysate at 20 µg

Lane 3:
MCF-7 (Human breast adenocarcinoma epithelial cell), Whole cell lysate at 20 µg

Lane 4:
U-87 MG (Human glioblastoma-astrocytoma epithelial cell), Whole cell lysate at 20 µg

Lane 5:
MDA-MB-435 (Human mammary gland ductal carcinoma melanocyte), Whole cell lysate at 20 µg

Lane 6:
T-47D (Human ductal breast epithelial tumor epithelial cell), Whole cell lysate at 20 µg

Lane 7:
Raji (Human Burkitt's lymphoma B lymphocyte), Whole cell lysate at 20 µg

Lane 8:
Ramos (Human Burkitt's lymphoma B lymphocyte), Whole cell lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Predicted band size: 43 kDa

Observed band size: 50 kDa

Exposure time: 60s



Western blot - Anti-p53 antibody [Y5] (ab32049)

P53 Western blot staining using rabbit Anti-p53 antibody

Blocking/Diluting buffer and concentration: 5% NFDM/TBST
Lane1: Mutant p53 cell lines

Lane2-3: Wildtype p53 cell lines

This antibody recognises both wild-type and mutant forms of human p53 and is not specific to any particular mutation. Expression levels of p53 can vary significantly between cell lines, with mutant forms often showing higher stability and abundance. In low-expressing wild-type cell lines, p53 levels may be increased using treatments such as camptothecin or irinotecan.

All lanes:
Western blot - Anti-p53 antibody [Y5] (ab32049) at 1/10000 dilution

Lane 1:
A431 (Human epidermoid carcinoma epithelial cell), Whole cell lysate at 20 µg

Lane 2:
A549 (Human lung carcinoma epithelial cell), Whole cell lysate at 20 µg


Lane 3:
MCF-7 (Human breast adenocarcinoma epithelial cell), Whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Predicted band size: 43 kDa

Exposure time: 180s



Western blot - Anti-p53 antibody [Y5] (ab32049)


P53 Western blot staining using rabbit Anti-p53 antibody

ab32049 was shown to react with TP53 in wild-type HAP1 cells in Western blot with loss of signal observed in a TP53 knockout cell line. Wild-type HAP1 and TP53 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab32049 overnight at 4 °C at a 1/500 dilution. Blots were incubated with secondary antibodies at 0.2 µg/mL before imaging.

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.
All lanes:
Western blot - Anti-p53 antibody [Y5] (ab32049) at 1/500 dilution

Lane 1:
Wild-type HAP1 lysate at 20 µg

Lane 2:
TP53 knock-out HAP1 lysate at 20 µg



Western blot - Anti-p53 antibody [Y5] (ab32049)

P53 Western blot staining using rabbit Anti-p53 antibody

False colour image of Western blot: Anti-p53 antibody [Y5] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab32049 was shown to bind specifically to p53. A band was observed at 50 kDa in wild-type HAP1 cell lysate with no signal observed at this size in tp53 knockout cell line. To generate this image, wild-type and tp53 knockout HAP1 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.
This antibody recognises both wild-type and mutant forms of human p53 and is not specific to any particular mutation. Expression levels of p53 can vary significantly between cell lines, with mutant forms often showing higher stability and abundance. In low-expressing wild-type cell lines, p53 levels may be increased using treatments such as camptothecin or irinotecan.

All lanes:
Western blot - Anti-p53 antibody [Y5] (ab32049) at 1/1000 dilution

Lane 1:
Saos-2 cell lysate at 20 µg

Lane 2:
A431 cell lysate at 20 µg

Lane 3:
Wild-type HAP1 cell lysate at 20 µg

Lane 4:
TP53 knockout HAP1 cell lysate at 20 µg

Lane 5:
MCF7 cell lysate at 20 µg

Lane 6:
HEK-293T cell lysate at 20 µg

Secondary


Lanes 1 - 6:
Western blot - Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) at 1/20000 dilution

Lanes 1 - 6:
Western blot - Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 43 kDa

Observed band size: 50 kDa



Western blot - Anti-p53 antibody [Y5] (ab32049)

P53 Western blot staining using rabbit Anti-p53 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

In Western blot, ab322465 was shown to bind specifically to TP53. Target of interest was observed at 35-53 kDa in wild-type HAP1 cell lysates (lane 1) with no signal observed at this size in TP53 knockout HAP1 cell line (lane 2).

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

In Western blot, Anti-p53 antibody [Y5] (ab32049) staining at 1/1000 dilution.

ab32049 recognises both wild-type and mutant forms of human p53 and is not specific to any particular mutation. Expression levels of p53 can vary significantly between cell lines, with mutant forms often showing higher stability and abundance. In low-expressing wild-type cell lines, p53 levels may be increased using treatments such as camptothecin or irinotecan.

All lanes:
Western blot - Anti-TP53 (phospho S376+S377+S392) antibody [RM1161] (ab322465) at 1/1000 dilution

Lane 1:
Wild-type HAP1 (human chronic myelogenous leukemia near-haploid cell) whole cell lysate at 20 µg


Lane 2:
TP53 knockout HAP1 whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 53 kDa, 35-48 kDa, 36 kDa

Exposure time: 15s



OI-RD Scanning - Anti-p53 antibody [Y5] (ab32049)

We have systematically measured KD (the equilibrium dissociation constant between the antibody and its antigen), of more than 840 recombinant antibodies to assess not only their individual KD values but also to see the average affinity of antibody. Based on the comparison with published literature values for mouse monoclonal antibodies, Recombinant antibodies appear to be on average 1-2 order of magnitude higher affinity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.