abcam

ab32059

Anti-cIAP2 antibody [E40]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	E40
Purification technique	Affinity purification Protein A
Specificity	This antibody is specific for human cIAP2. It does not cross-react with any other IAP family members.
Concentration	0.635 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Human
Dilution info	-

Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	**For unpurified use at 5 µg/mL.**

WB

Tested

Species	Human
Dilution info	1/50000
Notes	**For unpurified use at 1/1000.**

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/70
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/600
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information BIRC3

Function	Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, mitogenic kinase signaling and cell proliferation, as well as cell invasion and metastasis. Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and regulates both canonical and non-canonical NF-kappa-B signaling by acting in opposite directions: acts as a positive regulator of the canonical pathway and suppresses constitutive activation of non-canonical NF-kappa-B signaling. The target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, RIPK2, RIPK3, RIPK4, CASP3, CASP7, CASP8, IKBKE, TRAF1, and BCL10. Acts as an important regulator of innate immune signaling via regulation of Toll-like receptors (TLRs), Nodlike receptors (NLRs) and RIG-I like receptors (RLRs), collectively referred to as pattern recognition receptors (PRRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase-independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply

- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CIAP2 also called E3 ubiquitin-protein ligase and encoded by the BIRC3 gene functions to inhibit apoptosis. This protein belongs to the inhibitors of apoptosis (IAP) family and is around 69 kDa in mass. cIAP2 is widely expressed in various tissues including the epithelial and lymphoid cells. It operates mechanically by ubiquitinating substrates marking them for proteasomal degradation which is key in preventing cell death and maintaining cellular homeostasis.
Biological function summary	CIAP2 serves as a regulator in cell survival by functioning within signaling cascades. It participates in the formation of the TNF receptor complex which influences the NF-κB signaling pathway and inflammation response. By interacting with several proteins cIAP2 modulates downstream signaling effects that can influence immune responses and cellular proliferation helping cells adapt and survive stressful conditions.
Pathways	CIAP2 plays roles in both the NF-κB and the MAPK signaling pathways. Within these pathways cIAP2 interacts closely with proteins such as TRAF2 and RIP1. In the NF-κB pathway it helps in the regulation of inflammatory responses while in the MAPK pathway it contributes to cell growth and differentiation processes. These pathways highlight its function in regulating immune system responses and cell adaptability.
Associated diseases and disorders	CIAP2 maintains connections with cancer and chronic inflammation. Abnormal expression or mutations of cIAP2 can lead to tumorigenesis due to disrupted apoptotic control linking it closely to cancers like lymphomas and leukemias. Additionally cIAP2 influences inflammatory diseases like Crohn's disease through its interactions with proteins such as NOD2 which further impacts the disease pathology. Proper regulation of cIAP2 is essential in avoiding unwanted cell survival and excessive inflammation highlighting its significance in various pathological conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-cIAP2 antibody [E40] (ab32059)

CIAP2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-cIAP2 antibody


ab32059 (unpurified) at a 1/50 dilution staining cIAP2 in human normal spleen, using Immunohistochemistry, Paraffin Embedded Tissue.



Flow Cytometry (Intracellular) - Anti-cIAP2 antibody [E40] (ab32059)

CIAP2 Flow Cytometry (Intracellular) staining using rabbit Anti-cIAP2 antibody


Intracellular Flow Cytometry analysis of Raji (Human Burkitt's lymphoma B lymphocyte) cells labeling cIAP2 with purified ab32059 at 1/70 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluorr® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-cIAP2 antibody [E40] (ab32059)

CIAP2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-cIAP2 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast carcinoma tissue sections labeling cIAP2 with purified ab32059 at 1/600 dilution (1.12 µg/ml). Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunoprecipitation - Anti-cIAP2 antibody [E40] (ab32059)

ab32059 (purified) at 1/30 dilution (2ug) immunoprecipitating cIAP2 in Daudi whole cell lysates.
Lane 1: Daudi (Human Burkitt's lymphoma lymphoblast) whole cell lysates 10ug
Lane 2 (+): ab32059 & Daudi whole cell lysates
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab32059 in Daudi whole cell lysates
For western blotting, VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/1000 dilution.
Blocking and diluting buffer: 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-cIAP2 antibody [E40] (ab32059)


Predicted band size: 68 kDa



Immunoprecipitation - Anti-cIAP2 antibody [E40] (ab32059)

cIAP2 was immunoprecipitated using 0.5mg Daudi whole cell extract, 5µg of Rabbit polyclonal to cIAP2 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).
The antibody was incubated under agitation with Protein G beads for 10min, Daudi whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.
Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab32059 (unpurified).
Secondary: Mouse monoclonal [SB62a] Secondary Antibody to Rabbit IgG light chain (HRP) (ab99697).
Band: 72kDa; cIAP2
All lanes:
Immunoprecipitation - Anti-cIAP2 antibody [E40] (ab32059)

Predicted band size: 68 kDa



Western blot - Anti-cIAP2 antibody [E40] (ab32059)

All lanes:
Western blot - Anti-cIAP2 antibody [E40] (ab32059) at 1/50000 dilution


All lanes:
Raji (Human Burkitt's lymphoma B lymphocyte) whole cell lysates at 15 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 68 kDa

Observed band size: 72 kDa



Western blot - Anti-cIAP2 antibody [E40] (ab32059)

All lanes:
Western blot - Anti-cIAP2 antibody [E40] (ab32059) at 1/1000 dilution

All lanes:
Daudi cell lysate.

Predicted band size: 68 kDa

Observed band size: 72 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.