abcam

ab32117

Anti-HDAC2 antibody [Y461]

Anti-HDAC2 antibody [Y461] (ab32117) is a rabbit monoclonal antibody detecting HDAC2 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 100 publications
- Trusted since 2006

Recombinant | RabMAb | Advanced Validation | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	Y461
Purification technique	Affinity purification Protein A
Concentration	0.542 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ChIC/CUT&RUN-seq

Tested

Species	Human
Dilution info	5 µg
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	-
Notes	**Purified ab32117 at 1:1500 dilution.** Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	-
Notes	**Purified ab32117 at 1:1500 dilution.** Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IP

Tested

Species	Human

Dilution info	1/60
Notes	-

Expected

Species	Mouse
Dilution info	1/60
Notes	-

Species	Rat
Dilution info	1/60
Notes	-

WB

Tested

Species	Human
Dilution info	1/2000
Notes	-

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat
Dilution info	1/2000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	-

Species	Rat
Dilution info	1/250 - 1/500
Notes	-

Expected

Species	Mouse
Dilution info	1/250 - 1/500
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/60 - 1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse
Dilution info	1/60 - 1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Species	Rat
Dilution info	1/60 - 1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target data

See full target information HDAC2

Function Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:28497810). Histone deacetylation gives a tag for epigenetic repression and plays an important

role in transcriptional regulation, cell cycle progression and developmental events (By similarity). Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR (PubMed:12724404). Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (By similarity). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed:16428440, PubMed:28977666). Component of the SIN3B complex that represses transcription and counteracts the histone acetyltransferase activity of EP300 through the recognition H3K27ac marks by PHF12 and the activity of the histone deacetylase HDAC2 (PubMed:37137925). Also deacetylates non-histone targets: deacetylates TSHZ3, thereby regulating its transcriptional repressor activity (PubMed:19343227). May be involved in the transcriptional repression of circadian target genes, such as PER1, mediated by CRY1 through histone deacetylation (By similarity). Involved in MTA1-mediated transcriptional corepression of TFF1 and CDKN1A (PubMed:21965678). In addition to protein deacetylase activity, also acts as a protein-lysine deacylase by recognizing other acyl groups: catalyzes removal of (2E)-butenoyl (crotonyl), lactoyl (lactyl) and 2-hydroxyisobutanoyl (2-hydroxyisobutyryl) acyl groups from lysine residues, leading to protein decrotonylation, delactylation and de-2-hydroxyisobutyrylation, respectively (PubMed:28497810, PubMed:29192674, PubMed:35044827).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-HDAC2 antibody [Y461] (ab32117) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of HDAC2?
Anti-HDAC2 [Y461] (ab32117) specifically detects a band for HDAC2 (UniProt: Q92769) at a molecular weight of 55kDa.

Trusted by the scientific community
Anti-HDAC2 [Y461] (ab32117) was first used in a scientific publication in 2006 and has been cited over 100 times in

peer-reviewed journals.

Reviewed by scientists
Anti-HDAC2 [Y461] (ab32117) has over 10 independent reviews from customers.

Specificity confirmed
The specificity of Anti-HDAC2 antibody [Y461] (ab32117) has been confirmed by Western blot testing in HDAC2 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [Y461] also available for your convenience: ab32117, HRP - ab195851, Alexa Fluor® 488 - ab196471, Alexa Fluor® 647 - ab196518, PE - ab210827, Carrier free - ab213700, APC - ab319340, Alexa Fluor® 594 - ab319833, Alexa Fluor® 555 - ab319974, Alexa Fluor® 750 - ab321577

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	HDAC2 or Histone Deacetylase 2 is an important enzyme that plays a role in chromatin remodeling and gene expression regulation. It removes acetyl groups from histone tails allowing chromatin to condense and silencing gene expression. HDAC2 has a molecular weight of approximately 55 kDa. It is expressed in various tissues including brain heart and skeletal muscles indicating its importance in different physiological areas. HDAC2 proteins are often studied using methods like HDAC2 ELISA and HDAC2 anticuero in research settings.
Biological function summary	HDAC2 is involved in cell cycle regulation and differentiation. It is part of the co-repressor complex and interacts with other proteins such as Sin3 and NuRD to repress transcription. These interactions are vital for maintaining normal cellular functions and ensuring proper response to external signals. This protein is also connected to HEK293T cells which are a common model in scientific studies due to their robust expression of proteins like HDAC2.

Pathways	HDAC2 participates in the regulation of transcriptional activity and cellular stress response. It is an important player in the histone modification pathway interacting with proteins such as HDAC1 and transcriptional regulators like REST. HDAC2 is also linked with the Notch signaling pathway which is essential for cell fate decisions. These pathways highlight its role in both maintaining cellular homeostasis and adapting to changes in the cellular environment.
Associated diseases and disorders	HDAC2 has a significant connection to neurodegenerative diseases and various cancers. Alzheimer's disease for instance shows altered expression levels of HDAC2 affecting synaptic plasticity and memory formation. In cancer HDAC2 interacts with oncogenic proteins such as p53 influencing tumor progression and metastasis. Monitoring HDAC2 levels and activity could provide insights into disease mechanisms and potential therapeutic targets.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


17 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)


Immunohistochemical analysis of paraffin-embedded fixed (A) Wild-type HEK293T (human embryonic kidney epithelial cell) cell pellet. (B) HDAC2 knockout HEK293T (ab266590) cell pellet

staining HDAC2 with ab32117 at 1/10000 dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Counter-staining used was hematoxylin. Positive staining on (A) Wild-type HEK293T cell pellet, no staining on HDAC2 knockout HEK293T (ab266590) cell pellet. The section was incubated with ab32117 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [Y461] (ab32117)


Immunocytochemistry/ Immunofluorescence analysis of Wild-type HEK293T/HDAC2 KO HEK293T (HDAC2 knockout human embryonic kidney epithelial cell) (ab266590) cells labeling HDAC2 with ab32117 at 1/200 dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed secondary antibody at 1/1000 dilution (2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution (2.5 µg/ml). DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control. Confocal image showing nuclear staining in Parental HEK293T cell line. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC2 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast cancer tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry (Intracellular) - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Flow Cytometry (Intracellular) staining using rabbit Anti-HDAC2 antibody


Flow Cytometry analysis of K-562 (Human chronic myelogenous leukemia lymphoblast) cells labelling HDAC2 with Purified ab32117 at 1:20 dilution (5 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150081) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-HDAC2 antibody


Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling HDAC2 with Purified ab32117 at 1:50 dilution (2.1 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-HDAC2 antibody


ab32117 staining HDAC2 in wild-type HAP1 cells (top panel) and HDAC2 knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab32117 at 1/250 dilution and ab195889 at 1/250 dilution (shown in pseudocolour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC2 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human testis tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunoprecipitation - Anti-HDAC2 antibody [Y461] (ab32117)

Purified ab32117 at 1:20 dilution (0.5µg) immunoprecipitating HDAC2 in HeLa whole cell lysate.
Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10 µg.
Lane 2 (+): ab32117 + HeLa whole cell lysate.
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab32127 in HeLa whole cell lysate.
VeriBlot for IP Detection Reagent (HRP)(ab131366) (1:5000 dilution) was used for Western blotting.
Blocking Buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM/TBST.
Observed band size: kDa
All lanes:
Immunoprecipitation - Anti-HDAC2 antibody [Y461] (ab32117)


Predicted band size: 55 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC2 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat testis tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC2 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse testis tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-HDAC2 antibody [Y461] (ab32117)

HDAC2 Western blot staining using rabbit Anti-HDAC2 antibody

All lanes:
Western blot - Anti-HDAC2 antibody [Y461] (ab32117) at 1/10000 dilution

Lane 1:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg


Lane 2:
Mouse brain lysate at 20 µg

Lane 3:
Rat brain lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 55 kDa



Western blot - Anti-HDAC2 antibody [Y461] (ab32117)


HDAC2 Western blot staining using rabbit Anti-HDAC2 antibody

Lane 1: Wild type HAP1 whole cell lysate (20 µg)
Lane 2: HDAC2 knockout HAP1 whole cell lysate (20 µg)
Lane 3: A431 whole cell lysate (20 µg)
Lane 4: Hela whole cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab32117 observed at 60 kDa. Red - loading control, ab8245, observed at 37 kDa.

ab32117 was shown to specifically react with HDAC2 when HDAC2 knockout samples were used. Wild-type and HDAC2 knockout samples were subjected to SDS-PAGE. ab32117 and ab8245 (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/2000 and 1/10000 respectively. Blots were developed with 800CW Goat anti Rabbit and 680CW Goat anti Mouse secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.


All lanes:
Western blot - Anti-HDAC2 antibody [Y461] (ab32117)

Predicted band size: 55 kDa




ChIC/CUT&RUN sequencing - Anti-HDAC2 antibody [Y461] (ab32117)

CUT&RUN profiling with HDAC2 antibody demonstrates robust genome-wide enrichment in wild-type (WT) cells, which is markedly diminished in HDAC2 knockout (KO) cells. Heatmaps of genome-wide signal flanking annotated transcription start sites (TSSs, +/- 2 kbp) display CUT&RUN data generated using the CUTANA™ CUT&RUN Kit (EpiCypher 14-1048) with HDAC2 antibody (Abcam ab32117, 0.5 µg). 500,000 HEK293T WT or KO (Abcam ab266590) cells were used per reaction. IgG antibody was included as a negative control to assess non-specific background. Libraries were prepared using the CUTANA™ CUT&RUN Library Prep Kit (EpiCypher 14-1001). Sequencing was performed with paired-end 50 bp reads, and data were processed on CUTANA™ Cloud (cloud.epicypher.com) by alignment to the hg38 genome. Heatmaps were generated using deepTools (Ramнrez et al., Nucleic Acids Res. 2014; PMID: 24799436). Row-linked data are ranked by intensity relative to HDAC2 WT, with red indicating high localized enrichment and blue denoting background. Validated antibodies show genome-wide enrichment above IgG background consistent with HDAC2 binding in WT cells and near complete loss of signal in KO cells.




ChIC/CUT&RUN sequencing - Anti-HDAC2 antibody [Y461] (ab32117)

CUT&RUN profiling with HDAC2 antibody reveals the expected genomic enrichment pattern in wild-type (WT) cells, which is substantially reduced in HDAC2 knockout (KO) cells. Representative genome browser tracks show CUT&RUN data generated using the CUTANA™ CUT&RUN Kit (EpiCypher 14-1048) with HDAC2 antibody (Abcam ab32117, 0.5 µg). 500,000 HEK293T WT or KO (Abcam ab266590) cells were used per reaction. IgG, H3K4me3, and H3K27me3 antibodies were included as controls to assess non-specific background, active promoters, and repressed chromatin, respectively. Libraries were prepared using the CUTANA™ CUT&RUN Library Prep Kit (EpiCypher 14-1001). Sequencing was performed with paired-end 50 bp reads, and data were processed on CUTANA™ Cloud (cloud.epicypher.com) by alignment to the hg38 genome. Images were generated using Integrative Genomics Viewer (IGV, Broad Institute).



ChIC/CUT&RUN sequencing - Anti-HDAC2 antibody [Y461] (ab32117)

ChIC/CUT&RUN was performed using a pAG-MNAse at a final concentration of 700 ng/mL, 2 x 10^5 K-562 (Human chronic myelogenous leukemia lymphoblast) cells and 5 µg of ab32117 [Y461] . The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown. Additional screenshots of mapped reads can be found in the Protocol booklet in the Product Protocol section. The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Western blot - Anti-HDAC2 antibody [Y461] (ab32117)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-HDAC2 antibody [Y461] (ab32117) at 1/1000 dilution

Lane 1:
GH3 (rat pituitary epithelial cell) whole cell lysate at 20 µg

Lane 2:
L6 (rat skeletal muscle myoblast) whole cell lysate at 20 µg

Lane 3:
C6 (rat glial tumor glial cell) whole cell lysate at 20 µg

Lane 4:
AR42J (rat pancreatic tumor epithelial cell) whole cell lysate at 20 µg

Lane 5:
2.4G2 (rat B cell lymphoma B lymphocyte) whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 55 kDa

Observed band size: 60 kDa

Exposure time: 37s



Western blot - Anti-HDAC2 antibody [Y461] (ab32117)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-HDAC2 antibody [Y461] (ab32117) at 1/1000 dilution

Lane 1:
HT-22 (mouse hippocampal neuronal cell) whole cell lysate at 20 µg

Lane 2:
SW10 (mouse neuronal Schwann cell) whole cell lysate at 20 µg

Lane 3:
bEnd.3 (mouse brain endothelioma) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 55 kDa

Observed band size: 60 kDa

Exposure time: 26s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.