abcam

ab322304

FITC Anti-ICAM1 antibody [EPR22161-284]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	FITC
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR22161-284
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human

Dilution info	1/2500
Notes	-

Predicted

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information Icam1

Function

Cell adhesion molecule that functions as a receptor ligand of the signaling receptor ITGAL:ITGB2/LFA-1 (lymphocyte-function associated (LFA) molecule 1) ensuring leukocyte cell-cell adhesion, by providing a calibrated system to namely adjust T-cell killing to the antigen stimulation strength. Also functions as a ligand receptor of the signaling receptor ITGAM:ITGB2/MAC-1 ensuring adhesion between stimulated neutrophils and stimulated endothelial cells. During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation. Promotes cell aggregation in epithelial cells through interaction with MUC1.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images




Flow Cytometry (Intracellular) - FITC Anti-ICAM1 antibody [EPR22161-284] (ab322304)

ICAM1 Flow Cytometry (Intracellular) staining using rabbit Anti-ICAM1 antibody

Flow cytometry overlay histogram showing left Ramos positive cells and right negative HEK293 stained with ab322304 (red line). The cells were incubated in 1x PBS containing µg/ml human IgG and 10% normal goat serum followed by the antibody (ab322915) (1x 10^6 in 100 µl at 0.2 µg/ml (1/2500)) for 30min on ice.

Isotype control antibody (black line) was Rabbit IgG [EPR25A] FITC used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.



Flow Cytometry - FITC Anti-ICAM1 antibody [EPR22161-284] (ab322304)

ICAM1 Flow Cytometry staining using rabbit Anti-ICAM1 antibody

Flow cytometry overlay histogram showing left A20 positive cells and right negative L-929 stained with ab322304 (red line). The cells were incubated in 1x PBS containing 10%; normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab322304) (1x 10^6 in 100ul at 0.2ug/ml (1/2500)) for 30min on ice.

Isotype control antibody (black line) was Rabbit IgG [EPR25A] FITC used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.