abcam

ab322709

Anti-CXCL16 antibody [EPR29713-506]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR29713-506
Purification technique	Affinity purification Protein A
Concentration	0.496 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Dot

Tested

Species	Recombinant fragment - Human
Dilution info	1/1000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Recombinant fragment - Human
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Human, Recombinant fragment - Human
Dilution info	-
Notes	-

Target data

See full target information CXCL16

Function	Acts as a scavenger receptor on macrophages, which specifically binds to OxLDL (oxidized low density lipoprotein), suggesting that it may be involved in pathophysiology such as atherogenesis (By similarity). Induces a strong chemotactic response. Induces calcium mobilization. Binds to CXCR6/Bonzo.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

3 product images




Flow Cytometry - Anti-CXCL16 antibody [EPR29713-506] (ab322709)

CXCL16 Flow Cytometry staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CXCL16 antibody

Flow cytometric analysis of Human PBMC (human peripheral blood mononuclear cell) cells labelling CXCL16 with ab322709 at 1/50 dilution (1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Human PBMC were co-stained with CD11b conjugated BV421.
Gated on viable cell.




Flow Cytometry - Anti-CXCL16 antibody [EPR29713-506] (ab322709)

CXCL16 Flow Cytometry staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CXCL16 antibody

Flow cytometric analysis of Human PBMC (human peripheral blood mononuclear cell) cells labelling CXCL16 with ab322709 at 1/50 dilution (1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control.

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Human PBMC were co-stained with CD19 conjugated Alexa Fluor®647.
Gated on viable cell.



Dot Blot - Anti-CXCL16 antibody [EPR29713-506] (ab322709)

CXCL16 Dot Blot staining of Human CXCL16 protein using rabbit Anti-CXCL16 antibody

Dot blot analysis of CXCL16 using ab322709 at 1:1000 (0.496 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1:100,000 dilution.

Lane1: Human CXCL16 protein

Exposure time: 180 seconds.

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

All lanes:
Dot Blot - Anti-CXCL16 antibody [EPR29713-506] (ab322709) at 1/1000 dilution

All lanes:
Human CXCL16 protein

Secondary

All lanes:
Dot Blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.