abcam

ab323754

PE Anti-Podoplanin antibody [EPR27193-535]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	PE
Excitation/Emission	Ex: 480;565nm, Em: 578nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR27193-535
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	1/5000

Notes	-

ICC/IF

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information Pdpn

Function

Mediates effects on cell migration and adhesion through its different partners. During development plays a role in blood and lymphatic vessels separation by binding CLEC1B, triggering CLEC1B activation in platelets and leading to platelet activation and/or aggregation (PubMed:14522983, PubMed:15231832, PubMed:17616532, PubMed:20110424). Interaction with CD9, on the contrary, attenuates platelet aggregation and pulmonary metastasis induced by PDPN. Mediates effects on cell migration and adhesion through its different partners. Through MSN or EZR interaction promotes epithelial-mesenchymal transition (EMT) leading to ERZ phosphorylation and triggering RHOA activation leading to cell migration increase and invasiveness. Interaction with CD44 promotes directional cell migration in epithelial and tumor cells (By similarity). In lymph nodes (LNs), controls fibroblastic reticular cells (FRCs) adhesion to the extracellular matrix (ECM) and contraction of the actomyosin by maintaining ERM proteins (EZR; MSN and RDX) and MYL9 activation through association with unknown transmembrane proteins. Engagement of CLEC1B by PDPN promotes FRCs relaxation by blocking lateral membrane interactions leading to reduction of ERM proteins (EZR; MSN and RDX) and MYL9 activation (PubMed:25347465). Through binding with LGALS8 may participate in connection of the lymphatic endothelium to the surrounding extracellular matrix (By similarity). In keratinocytes, induces changes in cell morphology showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion (PubMed:10574709). Controls invadopodia stability and maturation leading to efficient degradation of the extracellular matrix (ECM) in tumor cells through modulation of RHOC activity in order to activate ROCK1/ROCK2 and LIMK1/LIMK2 and inactivation of CFL1 (By similarity). Required for normal lung cell proliferation and alveolus formation at birth (PubMed:12654292). Does not function as a water channel or as a regulator of aquaporin-type water channels (By similarity). Does not have any effect on folic acid or amino acid transport (PubMed:12032185).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Podoplanin also known as gp36 and PDPN is a small transmembrane protein weighing approximately 36 kDa. It mainly expresses in lymphatic endothelial cells podocytes and various other tissues like the lung and kidney. It functions as a marker in mesothelial cells and is noticeable in histochemical studies such as mesothelioma using podoplanin staining in immunohistochemistry (IHC). Podoplanin emerges as an important tool in identifying lymphatic vessels and has significant roles in the development of certain tissues.
Biological function summary	The protein participates in maintaining the integrity of cell structures and lymphangiogenesis. This protein does not form part of a larger complex but actively interacts with different molecules. In lymphoid tissues podoplanin contributes to the proper formation of lymphatic channels and assists in platelet aggregation. These actions are pivotal in wound healing and protecting the epithelial cell layers in multiple organs.

Pathways	Podoplanin impacts the platelet activation pathway and the Lymphatic Vessel Development pathway. Its interaction with CLEC-2 (C-type lectin-like receptor 2) triggers downstream signaling leading to changes in platelet morphology and actin cytoskeleton reorganization. PodoplaninŸÏs activation of these pathways links it to key processes that include cell migration and tissue homeostasis TGF-beta interaction also represents a significant relationship within these pathways.
Associated diseases and disorders	Podoplanin has correlations with conditions like cancer specifically mesothelioma and squamous cell carcinoma. Dysregulation of podoplanin and its pathways may contribute to tumor progression and metastasis. In mesothelioma podoplanin interacts with other proteins like E-cadherin influencing cancer cell migration and adhesion. Its role in disease states highlights the importance of podoplanin IHC as a diagnostic tool aiding in the better understanding and identification of disease pathology.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images




Flow Cytometry - PE Anti-Podoplanin antibody [EPR27193-535] (ab323754)

Podoplanin Flow Cytometry staining using rabbit Anti-Podoplanin antibody

Flow cytometric analysis of Hepa1-6 (mouse hepatoma epithelial cell, Left) / C2C12 (mouse myoblast, Right) cells labelling Podoplanin with ab323754 at 1/5000 dilution (0.01ug) (Red) compared with a Rabbit IgG monoclonal [EPR25A] - Isotype Control (Phycoerythrin) (ab209478) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Gated on viable cells.
Low expression: Hepa1-6.



Flow Cytometry - PE Anti-Podoplanin antibody [EPR27193-535] (ab323754)

Podoplanin Flow Cytometry staining using rabbit Anti-Podoplanin antibody

Flow cytometric analysis of RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage, Left) / NIH/3T3 (mouse embryonic fibroblast, Right) cells labelling Podoplanin with ab323754 at 1/5000 dilution (0.01ug) (Red) compared with a Rabbit IgG monoclonal [EPR25A] - Isotype Control (Phycoerythrin) (ab209478) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Gated on viable cells.
Low expression: RAW 264.7.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.