abcam

ab32420

Anti-ATM antibody [Y170]

Anti-ATM antibody [Y170] (ab32420) is a rabbit monoclonal antibody detecting ATM in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 140 publications
- Trusted since 2006

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	Y170
Purification technique	Affinity purification Protein A
Concentration	0.141 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Not recommended

Species	Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/20
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information ATM

Function

Serine/threonine protein kinase which activates checkpoint signaling upon double strand breaks (DSBs), apoptosis and genotoxic stresses such as ionizing ultraviolet A light (UVA), thereby acting as a DNA damage sensor (PubMed:10550055, PubMed:10839545, PubMed:10910365, PubMed:12556884, PubMed:14871926, PubMed:15064416, PubMed:15448695, PubMed:15456891, PubMed:15790808, PubMed:15916964, PubMed:17923702, PubMed:21757780, PubMed:24534091, PubMed:35076389, PubMed:9733514). Recognizes the substrate consensus sequence [ST]-Q (PubMed:10550055, PubMed:10839545, PubMed:10910365, PubMed:12556884, PubMed:14871926, PubMed:15448695, PubMed:15456891, PubMed:15916964, PubMed:17923702, PubMed:24534091, PubMed:9733514). Phosphorylates 'Ser-139' of histone variant H2AX at double strand breaks (DSBs), thereby regulating DNA damage response mechanism (By similarity). Also plays a role in pre-B cell allelic exclusion, a process leading to expression of a single immunoglobulin heavy chain allele to enforce clonality and monospecific recognition by the B-cell antigen receptor (BCR) expressed on individual B-lymphocytes. After the introduction of DNA breaks by the RAG complex on one immunoglobulin allele, acts by mediating a repositioning of the second allele to pericentromeric heterochromatin, preventing accessibility to the RAG complex and recombination of the second allele. Also involved in signal transduction and cell cycle control. May function as a tumor suppressor. Necessary for activation of ABL1 and SAPK. Phosphorylates DYRK2, CHEK2, p53/TP53, FBXW7, FANCD2, NFKBIA, BRCA1, CREBBP/CBP, RBBP8/CTIP, FBXO46, MRE11, nibrin (NBN), RAD50, RAD17, PELI1, TERF1, UFL1, RAD9, UBQLN4 and DCLRE1C (PubMed:10550055, PubMed:10766245, PubMed:10802669, PubMed:10839545, PubMed:10910365, PubMed:10973490, PubMed:11375976, PubMed:12086603, PubMed:15456891, PubMed:19965871, PubMed:21757780, PubMed:24534091, PubMed:26240375, PubMed:26774286, PubMed:30171069, PubMed:30612738, PubMed:30886146, PubMed:30952868, PubMed:38128537, PubMed:9733515, PubMed:9843217). May play a role in vesicle and/or protein transport. Could play a role in T-cell development, gonad and neurological function. Plays a role in replication-dependent histone mRNA degradation. Binds DNA ends. Phosphorylation of DYRK2 in nucleus in response to genotoxic stress prevents its MDM2-mediated ubiquitination and subsequent proteasome degradation (PubMed:19965871). Phosphorylates ATF2 which stimulates its function in DNA damage response (PubMed:15916964). Phosphorylates ERCC6 which is essential for its chromatin remodeling activity at DNA double-strand breaks (PubMed:29203878). Phosphorylates TTC5/STRAP at 'Ser-203' in the cytoplasm in response to DNA damage, which promotes TTC5/STRAP nuclear localization (PubMed:15448695). Also involved in pexophagy by mediating phosphorylation of PEX5: translocated to peroxisomes in response to reactive oxygen species (ROS), and catalyzes phosphorylation of PEX5, promoting PEX5 ubiquitination and induction of pexophagy (PubMed:26344566).

Storage

Shipped at conditions Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-ATM antibody [Y170] (ab32420) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human samples.

What is the molecular weight of ATM?
Anti-ATM [Y170] (ab32420) specifically detects a band for ATM (UniProt: Q13315) at a molecular weight of 350kDa.

Trusted by the scientific community
Anti-ATM [Y170] (ab32420) was first used in a scientific publication in 2006 and has been cited over 140 times in peer-reviewed journals.

Reviewed by scientists
Anti-ATM [Y170] (ab32420) has over 5 independent reviews from customers.

Specificity confirmed
The specificity of Anti-ATM antibody [Y170] (ab32420) has been confirmed by Western blot testing in ATM Knockout MCF7 cell line, ab282630.

Other related products
We have a range of other formats of antibody clone [Y170] also available for your convenience: ab32420, Alexa Fluor® 488 - ab208118, Carrier free - ab216617, PE - ab224947, APC - ab310856, Alexa Fluor® 647 - ab311093, Alexa Fluor® 594 - ab311698, Alexa Fluor® 568 - ab312973, Alexa Fluor® 555 - ab313182, Alexa Fluor® 750 - ab321602

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	ATM also known as Ataxia Telangiectasia Mutated is a protein kinase with a molecular weight of approximately 370 kDa. ATM protein primarily resides in the cell nucleus and functions as a critical regulator of the cell cycle. It plays a significant role in the detection of DNA damage and initiation of repair processes. As part of its mechanical functions ATM phosphorylates serine and threonine residues on various substrates most notably in response to double-strand breaks in DNA. This activity is important for maintaining genomic stability.
Biological function summary	ATM acts as a coordinator in cellular response to DNA damage highly interacting with multiple components of the DNA repair machinery. It forms a complex with proteins like NBS1 and MRN complex facilitating repair by recruiting and activating other proteins involved in homologous recombination and non-homologous end joining pathways. ATM also modulates p53 activity a primary response factor in cellular stress management linking ATM to control of cell cycle arrest and apoptosis. This positions ATM as an integral part of maintaining cellular integrity in face of genomic insult.
Pathways	ATM integrates neatly within the DNA damage response and cell cycle control pathways. ATM's operative relationship with the MRN complex and its role in the PI3K-related protein kinase family helps initiate appropriate repair processes upon DNA damage detection. Additionally ATM regulates the activity of proteins such as Chk2 which further propagates signals to p53 influencing decisions between cell cycle arrest and apoptosis. These interactions link ATM closely to essential processes like DNA repair and cell survival highlighting its role in genomic maintenance.
Associated diseases and disorders	ATM mutations or dysregulation leads to Ataxia Telangiectasia an autosomal recessive disorder characterized by neurodegeneration immune deficiencies and cancer predisposition. ATM dysfunction also connects to cancer development particularly breast cancer where it transmits signals involving BRCA1 contributing to DNA repair through homologous recombination. Understanding ATM dynamics and related pathways has important implications for developing therapeutic strategies to manage or mitigate effects associated with its dysfunction.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


12 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATM antibody [Y170] (ab32420)

ATM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ATM antibody


Formaldehyde-fixed human serous ovarian tumor tissue stained for ATM using ab32420 at 1/50 dilution in immunohistochemical analysis.



Flow Cytometry (Intracellular) - Anti-ATM antibody [Y170] (ab32420)

ATM Flow Cytometry (Intracellular) staining using rabbit Anti-ATM antibody


Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labelling ATM with purified ab32420 at 1/20 (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. An Alexa Fluorr® 488-conjugated goat anti-rabbit IgG (1/2000) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATM antibody [Y170] (ab32420)

ATM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ATM antibody


Formaldehyde-fixed human colon tissue stained for ATM using ab32420 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATM antibody [Y170] (ab32420)

ATM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human breast carcinoma using rabbit Anti-ATM antibody


Immunohistochemical analysis of formalin fixed paraffin embedded human breast carcinoma labelling ATM with ab32420 at a concentration of 0.1µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with a OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was performed with DISCOVERY cell conditioning solution (CC1) 100°C, pH8.5 for 32mins. ab32420 anti-ATM antibody [Y170] was incubated for 16mins at 37°C. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.
Customers are encouraged to optimise antigen retrieval conditions, antibody concentration, incubation times and temperature for best results in their own IHC assay workflow (automated and manual).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATM antibody [Y170] (ab32420)

ATM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ATM antibody


Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue labeling ATM with ab32420 at 1:100 dilution. Tissue underwent antigen retrieval using Tris/EDTA Buffer (pH9.0). The section was counterstained with haematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATM antibody [Y170] (ab32420)

ATM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ATM antibody


Immunohistochemical analysis of paraffin-embedded Human normal breast tissue labeling ATM with ab32420 at 1:100 dilution. Tissue underwent antigen retrieval using Tris/EDTA Buffer (pH9.0). The section was counterstained with haematoxylin.




Immunocytochemistry/ Immunofluorescence - Anti-ATM antibody [Y170] (ab32420)

ATM Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ATM antibody

Immunocytochemistry/ Immunofluorescence analysis of HeLa cells labeling ATM with ab32420 at 1/500. Goat anti rabbit IgG(Alexa Fluor® 488), ab150077 at 1/1000 was used as the secondary antibody. Cells were fixed with 4% Paraformaldehyde and permeabilised with 0.1% tritonX-100. Nulei were counterstained blue with DAPI.






Immunocytochemistry/ Immunofluorescence - Anti-ATM antibody [Y170] (ab32420)

ATM Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ATM antibody

Immunocytochemistry/ Immunofluorescence analysis of HepG2 cells labeling ATM with ab32420 at 1/500. Goat anti rabbit IgG(Alexa Fluor® 488), ab150077 at 1/1000 was used as the secondary antibody. Cells were fixed with 4% Paraformaldehyde and permeabilised with 0.1% tritonX-100. Nulei were counterstained blue with DAPI.



Western blot - Anti-ATM antibody [Y170] (ab32420)

ATM Western blot staining using rabbit Anti-ATM antibody

Western blot: Anti-ATM antibody [Y170] (ab32420) staining at 1/1000 dilution, shown in green; Mouse anti-CANX [CANX/1543] (ab238078) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab32420 was shown to bind specifically to ATM. A band was observed at 350 kDa in wild-type MCF7 cell lysates with a reduction in signal observed at this size in ATM heterozygous knockout cell line ab282630. To generate this image, wild-type and ATM heterozygous knockout MCF7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-ATM antibody [Y170] (ab32420) at 1/1000 dilution


Lane 1:
Wild-type MCF7 cell lysate at 20 µg

Lane 2:
ATM knockout MCF7 cell lysate at 20 µg

Lane 3:
Wild-type A549 cell lysate at 20 µg

Lane 4:
ATM knockout A549 ab283811 cell lysate at 20 µg

Observed band size: 350 kDa



Western blot - Anti-ATM antibody [Y170] (ab32420)

ATM Western blot staining using rabbit Anti-ATM antibody

False colour image of Western blot: Anti-ATM antibody [Y170] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (ab7291) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab32420 was shown to bind specifically to ATM. A band was observed at 350 kDa in wild-type A549 cell lysates with no signal observed at this size in ATM knockout cell line ab276095 (knockout cell lysate ab283834). To generate this image, wild-type and ATM knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5% milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.
All lanes:
Western blot - Anti-ATM antibody [Y170] (ab32420) at 1/1000 dilution

Lane 1:
Wild-type A549 cell lysate at 20 µg

Lane 2:
ATM knockout A549 cell lysate at 20 µg


Lane 3:
HEK-293 cell lysate at 20 µg

Lane 4:
U-2 OS cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 351 kDa

Observed band size: 350 kDa



Western blot - Anti-ATM antibody [Y170] (ab32420)

All lanes:
Western blot - Anti-ATM antibody [Y170] (ab32420) at 1/3000 dilution

Lane 1:
HeLa cell lysate at 10 µg

Lane 2:
HEK293 cell lysate at 10 µg

Secondary

All lanes:
HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size: 351 kDa

Observed band size: 370 kDa




Image from Adeyemi R.O. et al PLoS Pathog. 2010 Oct 7;6(10):e1001141. doi: 10.1371/journal.ppat.1001141.

Immunocytochemistry/ Immunofluorescence - Anti-ATM antibody [Y170] (ab32420)

DNA repair proteins accumulate at MVM APAR bodies
Repair proteins accumulate at APAR bodies. NB324K cells were infected with MVMp (MOI of 10) for 16 hr before being fixed and processed for immunofluorescence. Cells were stained with the indicated antibodies to mark DDR repair proteins. APAR bodies were detected with antibodies to NS1. Nuclei were stained with DAPI. All images were captured using an objective of 63×.

Cells were fixed with 4% paraformaldehyde for 15 minutes and permeabilized with 0.5% Triton X-100 in PBS for 15 minutes.

(Image shows the right-hand panel of Figure 2A)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.