abcam

ab324717

Anti-TIGIT antibody [EPR28147-69]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR28147-69
Purification technique	Affinity purification Protein A
Concentration	0.479 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Target data

See full target information Tigit

Function	Inhibitory receptor that plays a role in the modulation of immune responses. Suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells. Upon binding to its ligands PVR/CD155 or NECTIN2/CD112, which are expressed on antigen-presenting cells, sends inhibitory signals to the T-cell or NK cell. Mechanistically, interaction with ligand leads to phosphorylation of the cytoplasmic tail by Src family tyrosine kinases such as FYN or LCK, allowing subsequent binding to adapter GRB2 and SHIP1/INPP5D. In turn, inhibits PI3K and MAPK signaling cascades. In addition, associates with beta-arrestin-2/ARRB2 to recruit SHIP1/INPP5D that suppresses autoubiquitination of TRAF6 and subsequently inhibits NF-kappa-B signaling pathway. Also acts as a receptor for NECTIN4 to inhibit NK cell cytotoxicity.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

3 product images




Flow Cytometry - Anti-TIGIT antibody [EPR28147-69] (ab324717)

TIGIT Flow Cytometry staining of Mouse spleen cells were treated with 5ug/ml of immobilized anti-CD3 and anti-CD28 for 24 hours (Lower left and right) using rabbit Anti-TIGIT antibody

Flow cytometric analysis of Mouse spleen cells were treated with 5ug/ml of immobilized anti-CD3 and anti-CD28 for 24 hours (Lower left and right) / Untreated control (Upper left and right) cells labelling TIGIT with ab324717 at 1/500 dilution (0.1ug) / Upper right and Lower right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with CD3 conjugated to Alexa Fluor® 647.




Flow Cytometry - Anti-TIGIT antibody [EPR28147-69] (ab324717)

TIGIT Flow Cytometry staining of Mouse spleen cells were treated with 5ug/ml of immobilized anti-CD3 and anti-CD28 for 24 hours (Lower left and right) using rabbit Anti-TIGIT antibody

Flow cytometric analysis of Mouse spleen cells were treated with 5ug/ml of immobilized anti-CD3 and anti-CD28 for 24 hours (Lower left and right) / Untreated control (Upper left and right) cells labelling TIGIT with ab324717 at 1/500 dilution (0.1ug) / Upper right and Lower right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with CD25 conjugated to Brilliant Violet 421.



Flow Cytometry - Anti-TIGIT antibody [EPR28147-69] (ab324717)

TIGIT Flow Cytometry staining of Mouse spleen cells were treated with 5ug/ml of immobilized anti-CD3 and anti-CD28 for 24 hours (Lower left and right) using rabbit Anti-TIGIT antibody

Flow cytometric analysis of Mouse spleen cells were treated with 5ug/ml of immobilized anti-CD3 and anti-CD28 for 24 hours (Lower left and right) / Untreated control (Upper left and right) cells labelling TIGIT with ab324717 at 1/500 dilution (0.1ug) / Upper right and Lower right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with CD19 conjugated to PE/Cy7.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.