abcam

ab32505

# Anti-AKT3 + AKT2 + AKT1 antibody [Y89]

Lab Essentials | Recombinant | RabMAb | 20ul selling size

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | Y89 |
| **Purification technique** | Affinity purification Protein A |
| **Specificity** | This product reacts with AKT1, AKT2 and AKT3. |
| **Concentration** | 0.102 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

## IP

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/50 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow |
| **Dilution info** | - |
| **Notes** | - |

## WB

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/2000 - 1/10000 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow |
| **Dilution info** | - |
| **Notes** | - |

## ICC/IF

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100 - 1/250 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow |
| **Dilution info** | - |
| **Notes** | - |

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/20 |
| **Notes** | ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow |
| **Dilution info** | - |
| **Notes** | - |

## IHC-P

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100 |
| **Notes** | Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow |
| **Dilution info** | - |
| **Notes** | - |

# Target data

See full target information AKT1

| | |
|---|---|
| **Function** | AKT1 is one of 3 closely related serine/threonine-protein kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which regulate many processes including metabolism, proliferation, cell survival, growth and angiogenesis (PubMed:11882383, PubMed:15526160, PubMed:15861136, PubMed:21432781, PubMed:21620960, PubMed:31204173). This is mediated through serine and/or threonine phosphorylation of a range of downstream substrates (PubMed:11882383, PubMed:15526160, PubMed:21432781, PubMed:21620960, PubMed:29343641, PubMed:31204173). Over 100 substrate candidates have been reported so far, but for most of them, no isoform specificity has been reported |

(PubMed:11882383, PubMed:15526160, PubMed:21432781, PubMed:21620960). AKT is responsible of the regulation of glucose uptake by mediating insulin-induced translocation of the SLC2A4/GLUT4 glucose transporter to the cell surface (By similarity). Phosphorylation of PTPN1 at 'Ser-50' negatively modulates its phosphatase activity preventing dephosphorylation of the insulin receptor and the attenuation of insulin signaling (By similarity). Phosphorylation of TBC1D4 triggers the binding of this effector to inhibitory 14-3-3 proteins, which is required for insulin-stimulated glucose transport (PubMed:11994271). AKT also regulates the storage of glucose in the form of glycogen by phosphorylating GSK3A at 'Ser-21' and GSK3B at 'Ser-9', resulting in inhibition of its kinase activity (By similarity). Phosphorylation of GSK3 isoforms by AKT is also thought to be one mechanism by which cell proliferation is driven (By similarity). AKT also regulates cell survival via the phosphorylation of MAP3K5 (apoptosis signal-related kinase) (PubMed:11154276). Phosphorylation of 'Ser-83' decreases MAP3K5 kinase activity stimulated by oxidative stress and thereby prevents apoptosis (PubMed:11154276). AKT mediates insulin-stimulated protein synthesis by phosphorylating TSC2 at 'Ser-939' and 'Thr-1462', thereby activating the mTORC1 signaling pathway, and leading to both phosphorylation of 4E-BP1 and in activation of RPS6KB1 (PubMed:12150915, PubMed:12172553). Also regulates the mTORC1 signaling pathway by catalyzing phosphorylation of CASTOR1 and DEPDC5 (PubMed:31548394, PubMed:33594058). AKT plays a role as key modulator of the AKT-mTOR signaling pathway controlling the tempo of the process of newborn neurons integration during adult neurogenesis, including correct neuron positioning, dendritic development and synapse formation (By similarity). Part of a positive feedback loop of mTORC2 signaling by mediating phosphorylation of MAPKAP1/SIN1, promoting mTORC2 activation (By similarity). AKT is involved in the phosphorylation of members of the FOXO factors (Forkhead family of transcription factors), leading to binding of 14-3-3 proteins and cytoplasmic localization (PubMed:10358075). In particular, FOXO1 is phosphorylated at 'Thr-24', 'Ser-256' and 'Ser-319' (PubMed:10358075). FOXO3 and FOXO4 are phosphorylated on equivalent sites (PubMed:10358075). AKT has an important role in the regulation of NF-kappa-B-dependent gene transcription and positively regulates the activity of CREB1 (cyclic AMP (cAMP)-response element binding protein) (PubMed:9829964). The phosphorylation of CREB1 induces the binding of accessory proteins that are necessary for the transcription of pro-survival genes such as BCL2 and MCL1 (PubMed:9829964). AKT phosphorylates 'Ser-454' on ATP citrate lyase (ACLY), thereby potentially regulating ACLY activity and fatty acid synthesis (By similarity). Activates the 3B isoform of cyclic nucleotide phosphodiesterase (PDE3B) via phosphorylation of 'Ser-273', resulting in reduced cyclic AMP levels and inhibition of lipolysis (By similarity). Phosphorylates PIKFYVE on 'Ser-318', which results in increased PI(3)P-5 activity (By similarity). The Rho GTPase-activating protein DLC1 is another substrate and its phosphorylation is implicated in the regulation cell proliferation and cell growth (By similarity). Signals downstream of phosphatidylinositol 3-kinase (PI(3)K) to mediate the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor 1 (IGF1) (PubMed:12176338, PubMed:12964941). AKT mediates the antiapoptotic effects of IGF1 (By similarity). Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly (PubMed:19934221). May be involved in the regulation of the placental development (By similarity). Phosphorylates STK4/MST1 at 'Thr-120' and 'Thr-387' leading to inhibition of its: kinase activity, nuclear translocation, autophosphorylation and ability to phosphorylate FOXO3 (PubMed:17726016). Phosphorylates STK3/MST2 at 'Thr-117' and 'Thr-384' leading to inhibition of its: cleavage, kinase activity, autophosphorylation at Thr-180,

binding to RASSF1 and nuclear translocation (PubMed:20086174). Phosphorylates SRPK2 and enhances its kinase activity towards SRSF2 and ACIN1 and promotes its nuclear translocation (PubMed:19592491). Phosphorylates RAF1 at 'Ser-259' and negatively regulates its activity (PubMed:10576742). Phosphorylation of BAD stimulates its pro-apoptotic activity (PubMed:10926925). Phosphorylates KAT6A at 'Thr-369' and this phosphorylation inhibits the interaction of KAT6A with PML and negatively regulates its acetylation activity towards p53/TP53 (PubMed:23431171). Phosphorylates palladin (PALLD), modulating cytoskeletal organization and cell motility (PubMed:20471940). Phosphorylates prohibitin (PHB), playing an important role in cell metabolism and proliferation (PubMed:18507042). Phosphorylates CDKN1A, for which phosphorylation at 'Thr-145' induces its release from CDK2 and cytoplasmic relocalization (PubMed:16982699). These recent findings indicate that the AKT1 isoform has a more specific role in cell motility and proliferation (PubMed:16139227). Phosphorylates CLK2 thereby controlling cell survival to ionizing radiation (PubMed:20682768). Phosphorylates PCK1 at 'Ser-90', reducing the binding affinity of PCK1 to oxaloacetate and changing PCK1 into an atypical protein kinase activity using GTP as donor (PubMed:32322062). Also acts as an activator of TMEM175 potassium channel activity in response to growth factors: forms the lysoK(GF) complex together with TMEM175 and acts by promoting TMEM175 channel activation, independently of its protein kinase activity (PubMed:32228865). Acts as a regulator of mitochondrial calcium uptake by mediating phosphorylation of MICU1 in the mitochondrial intermembrane space, impairing MICU1 maturation (PubMed:30504268). Acts as an inhibitor of tRNA methylation by mediating phosphorylation of the N-terminus of METTL1, thereby inhibiting METTL1 methyltransferase activity (PubMed:15861136). In response to LPAR1 receptor pathway activation, phosphorylates Rabin8/RAB3IP which alters its activity and phosphorylates WDR44 which induces WDR44 binding to Rab11, thereby switching Rab11 vesicular function from preciliary trafficking to endocytic recycling (PubMed:31204173).

## Storage

| | |
|---|---|
| **Shipped at conditions** | Conditional Ambient |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | -20°C |
| **Aliquoting information** | Upon delivery aliquot |
| **Storage information** | Avoid freeze / thaw cycle |

## Notes

**Patented technology**

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

## Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

| | |
|---|---|
| **Activity summary** | The protein kinase AKT also known as protein kinase B (PKB) includes three isoforms: AKT1 AKT2 and AKT3. These play a central role in cellular processes. AKT1 has a molecular weight of approximately 56 kDa AKT2 has around 55 kDa and AKT3 similarly weighs about 54 kDa. These isoforms of AKT are expressed in various tissues but AKT1 is abundantly found in most tissues AKT2 is often present in insulin-responsive tissues like skeletal muscle and adipose tissue and AKT3 is mainly in the brain. The activation of AKT involves phosphorylation which increases their kinase activity significantly. The isoforms have unique and overlapping functions offering distinct mechanical roles in cellular signaling. |
| **Biological function summary** | The AKT kinase family plays a major role in regulating vital cellular functions including metabolism cell proliferation survival and growth. It does so by catalyzing phosphorylation of a range of substrates within these pathways. AKT is often involved in complexes with other proteins to achieve these biological effects. It impacts the regulation of glycogen synthase kinase 3 (GSK3) involved in insulin response and mTOR which controls cell growth. These wide-ranging functions make AKT proteins essential in maintaining cellular homeostasis. |
| **Pathways** | AKT proteins are important components of the PI3K/AKT/mTOR signaling pathway and the insulin signaling pathway. Within these pathways they interact with and modulate activities of related proteins such as PI3K PDK1 and mTOR. This regulation helps control the balance between anabolic and catabolic processes influencing cell survival growth and metabolism. The importance of AKT kinases in these pathways marks them as key regulators of cellular responses to external and internal stimuli. |
| **Associated diseases and disorders** | AKT isoforms have strong connections to cancer and diabetes. Over-activation of AKT pathways frequently occurs in cancerous cells contributing to uncontrolled cell growth and survival. In diabetes AKT2 is particularly implicated due to its role in insulin signaling influencing how cells process glucose. Dysregulation of AKT pathways has been linked to the development and progression of these diseases where proteins like mTOR in cancer and IRS1 in diabetes show direct relation to AKT function and malfunctions. |

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**

We do not recommend this combination. It is not covered by our product promise.


---

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


## 7 product images



### Immunocytochemistry/ Immunofluorescence - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)

AKT3 + AKT2 + AKT1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-AKT3 + AKT2 + AKT1 antibody


ab32505 staining in SK-N-SH cells treated with alsterpaullone (ab141070), by ICC/IF. Decrease of AKT1 + AKT2 + AKT3 expression correlates with increased concentration of alsterpaullone, as described in literature.
The cells were incubated at 37°C for 6h in media containing different concentrations of ab141070 (alsterpaullone) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab32505 (1/200 dilution was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 anti-rabbit polyclonal antibody (ab96899) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.




### Flow Cytometry (Intracellular) - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)

AKT3 + AKT2 + AKT1 Flow Cytometry (Intracellular) staining using rabbit Anti-AKT3 + AKT2 + AKT1 antibody


Overlay histogram showing HeLa cells stained with ab32505 (red line). The cells were fixed with methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab32505, 1/20 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (ab96899) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit monoclonal IgG (1µg/1x$10^6$ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a slightly decreased signal in HeLa cells fixed with 4% paraformaldehyde (10 min)/permeabilized in 0.1% PBS-Tween used under the same conditions.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)


AKT3 + AKT2 + AKT1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-AKT3 + AKT2 + AKT1 antibody

Immunohistochemical analysis of paraffin-embedded prostate carcinoma using ab32505 at 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




## Immunoprecipitation - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)

Purified ab32505 at 1/50 dilution (2µg) immunoprecipitating AKT3+AKT2+AKT1 in MCF7 whole cell lysate.
Lane 1 (input): MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate 10µg
Lane 2 (+): ab32505 + MCF7 whole cell lysate.
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab32505 in MCF7 whole cell lysate.
VeriBlot for IP Detection Reagent (HRP) (ab131366) (1/1000 dilution) was used for Western blotting.
Blocking Buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM/TBST.
Observed band size: 59 kDa
All lanes:
Immunoprecipitation - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)





## Western blot - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)

AKT3 + AKT2 + AKT1 Western blot staining of MCF-7 cell lysate using rabbit Anti-AKT3 + AKT2 + AKT1 antibody

All lanes:
Western blot - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505) at 1/10000 dilution

All lanes:
MCF-7 cell lysate

Observed band size: 59 kDa




## Western blot - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)

AKT3 + AKT2 + AKT1 Western blot staining using rabbit Anti-AKT3 + AKT2 + AKT1 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505) at 1/2000 dilution

Lane 1:
293T cell lysate transfected with GFP tagged AKT2 at 10 µg


Lane 2:
293T cell lysate transfected with empty vector at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 82 kDa

Exposure time: 5s




## Western blot - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505)

AKT3 + AKT2 + AKT1 Western blot staining using rabbit Anti-AKT3 + AKT2 + AKT1 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-AKT3 + AKT2 + AKT1 antibody [Y89] (ab32505) at 1/10000 dilution

Lane 1:
293T cell lysate transfected with GFP tagged AKT1 at 10 µg

Lanes 2 and 4:
293T cell lysate transfected with empty vector at 10 µg

Lane 3:
293T cell lysate transfected with GFP tagged AKT3 at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 82 kDa

Exposure time: 8s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.