abcam

ab32516

Anti-AIF antibody [E20] - Mitochondrial Marker

Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516) is a rabbit monoclonal antibody detecting AIF in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, IHC-Fr, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 70 publications
- Trusted since 2006

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	E20
Concentration	0.086 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human

Dilution info	-
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Predicted

Species	Rat, Mouse
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Rat, Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Predicted

Species	Rat, Mouse
Dilution info	-
Notes	-

IHC-Fr

Expected

Species	Mouse, Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Rat, Mouse
Dilution info	-

Notes	-

Target data

See full target information AIFM1

Function	Functions both as NADH oxidoreductase and as regulator of apoptosis (PubMed:17094969, PubMed:20362274, PubMed:23217327, PubMed:33168626). In response to apoptotic stimuli, it is released from the mitochondrion intermembrane space into the cytosol and to the nucleus, where it functions as a proapoptotic factor in a caspase-independent pathway (PubMed:20362274). Release into the cytoplasm is mediated upon binding to poly-ADP-ribose chains (By similarity). The soluble form (AIFsol) found in the nucleus induces 'parthanatos' i.e. caspase-independent fragmentation of chromosomal DNA (PubMed:20362274). Binds to DNA in a sequence-independent manner (PubMed:27178839). Interacts with EIF3G, and thereby inhibits the EIF3 machinery and protein synthesis, and activates caspase-7 to amplify apoptosis (PubMed:17094969). Plays a critical role in caspase-independent, pyknotic cell death in hydrogen peroxide-exposed cells (PubMed:19418225). In contrast, participates in normal mitochondrial metabolism. Plays an important role in the regulation of respiratory chain biogenesis by interacting with CHCHD4 and controlling CHCHD4 mitochondrial import (PubMed:26004228). Isoform 4. Has NADH oxidoreductase activity. Does not induce nuclear apoptosis. Isoform 5. Pro-apoptotic isoform.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?

Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunohistochemistry (IHC-Fr), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat

samples.

What is the molecular weight of AIF?
Anti-AIF [E20] - Mitochondrial Marker (ab32516) specifically detects a band for AIF (UniProt: O95831) at a molecular weight of 67kDa.

Trusted by the scientific community
Anti-AIF [E20] - Mitochondrial Marker (ab32516) was first used in a scientific publication in 2006 and has been cited over 70 times in peer-reviewed journals.

Reviewed by scientists
Anti-AIF [E20] - Mitochondrial Marker (ab32516) has over 5 independent reviews from customers.

Specificity confirmed
The specificity of Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516) has been confirmed by Western blot testing in AIFM1 Knockout HEK-293 cells.

Other related products
We have a range of other formats of antibody clone [E20] also available for your convenience: ab32516, Alexa Fluor® 647 - ab196847, Biotin - ab197526, Carrier free - ab220215

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	AIF short for Apoptosis-Inducing Factor is a flavoprotein characterized by its mass of approximately 57 kDa. It resides in the mitochondria of cells where it plays a decisive role in apoptosis. Under normal conditions AIF remains in the mitochondrial intermembrane space but when cells receive apoptotic signals AIF translocates to the nucleus leading to chromatin condensation and large-scale DNA fragmentation. The protein is also known as AIFM1 and performs functions beyond apoptosis including regulation of reactive oxygen species. Often analyzed

	using tools like a mitochondrial AB marker researchers use AIF as an important part of understanding cell death mechanisms.
Biological function summary	AIF functions as an important element in mitochondrial-mediated apoptosis. Once released from mitochondria AIF triggers a caspase-independent pathway of apoptosis making it distinct from other pathways where caspases are central. While not part of a larger protein complex AIF interacts closely with several mitochondrial and nuclear elements to execute its functions. Researchers often monitor AIF activity using apoptosis ELISA kits which help in precise detection and quantification of this protein within cellular systems.
Pathways	The presence and activity of AIF align with the broader apoptotic pathway and mitochondrial respiration. AIF links significantly with other molecules like cytochrome c released during apoptosis although cytochrome c follows a caspase-dependent pathway. It also shows involvement in various cellular injury and stress response pathways reinforcing its role as a major player in cellular fate decisions and intrinsic apoptotic mechanisms.
Associated diseases and disorders	AIF holds relevance in neurodegenerative disorders such as Parkinson's disease and certain forms of cancer. Misregulation or mutations in AIF can lead to increased susceptibility to these conditions highlighting its importance in maintaining normal cellular and mitochondrial function. AIF also interacts with other proteins implicated in these disorders such as E2F1 in cancer which shares similar apoptotic regulatory roles. These interactions highlight the biological significance of AIF beyond its fundamental apoptosis function making it a potential target for therapeutic interventions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

AIF Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-AIF antibody


ab32516, at a 1/500 dilution, staining AIF in paraffin embedded human cervical carcinoma tissue by Immunohistochemistry.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

AIF Flow Cytometry (Intracellular) staining using rabbit Anti-AIF antibody


Overlay histogram showing K562 cells stained with ab32516 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab32516, 1/50 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (ab96899) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit monoclonal IgG (1µg/1x10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in K562 cells fixed with methanol (5 min)/permeabilized with 0.1% PBS-Tween used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

AIF Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-AIF antibody


ICC/IF image of ab32516 stained HeLa cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab32516, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Immunoprecipitation - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

AIF was immunoprecipitated using 0.5mg K562 whole cell extract, 5µg of Rabbit monoclonal to AIF and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).
The antibody was incubated under agitation with Protein G beads for 10min, K562 whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.
Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab32516.
Secondary: Mouse monoclonal [SB62a] Secondary Antibody to Rabbit IgG light chain (HRP) (ab99697).
Band: 67kDa: AIF
All lanes:
Immunoprecipitation - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

Predicted band size: 67 kDa



Western blot - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

AIF Western blot staining using rabbit Anti-AIF antibody

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

In Western blot, Anti-AIF antibody [E20] (ab32516) staining at 1/1000 dilution.

All lanes:
Western blot - Anti-CISD1/MitoNEET antibody [EPR29116-27] (ab318199) at 1/1000 dilution

Lane 1:
HeLa (human cervical adenocarcinoma epithelial cell) cytoplasmic fraction at 20 µg


Lane 2:
HeLa mitochondrial fraction at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Observed band size: 15 kDa, 36 kDa, 67 kDa

Exposure time: 10s



Western blot - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

AIF Western blot staining using rabbit Anti-AIF antibody

Lanes 1- 2: Merged signal (red and green). Green - ab32516 observed at 67 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (ab8245) observed at 37 kDa.
ab32516 was shown to react with AIF in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line ab266347 (knockout cell lysate ab256834) was used. Wild-type HEK-293T and AIFM1 knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab32516 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516) at 1/1000 dilution

Lane 1:
Wild-type HEK-293T cell lysate at 20 µg


Lane 2:
AIFM1 knockout HEK-293T cell lysate at 20 µg

Lane 2:
Western blot - Human AIFM1 (AIF) knockout HEK-293T cell line (ab266347)

Performed under reducing conditions.

Predicted band size: 67 kDa

Observed band size: 67 kDa



Western blot - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516)

All lanes:
Western blot - Anti-AIF antibody [E20] - Mitochondrial Marker (ab32516) at 1/1000 dilution

All lanes:
K562 cell lysate

Predicted band size: 67 kDa

Observed band size: 67 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.