abcam

ab325180

Anti-CD101 antibody [EPR28974-15]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR28974-15
Purification technique	Affinity purification Protein A
Concentration	0.497 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Dot

Tested

Species	Synthetic peptide - Human
Dilution info	1/1000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Synthetic peptide - Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Synthetic peptide - Human

Dilution info	-
Notes	-

Target data

See full target information CD101

Function	Plays a role as inhibitor of T-cells proliferation induced by CD3. Inhibits expression of IL2RA on activated T-cells and secretion of IL2. Inhibits tyrosine kinases that are required for IL2 production and cellular proliferation. Inhibits phospholipase C-gamma-1/PLCG1 phosphorylation and subsequent CD3-induced changes in intracellular free calcium. Prevents nuclear translocation of nuclear factor of activated T-cell to the nucleus. Plays a role in the inhibition of T-cell proliferation via IL10 secretion by cutaneous dendritic cells. May be a marker of CD4(+) CD56(+) leukemic tumor cells.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

6 product images




Flow Cytometry - Anti-CD101 antibody [EPR28974-15] (ab325180)

CD101 Flow Cytometry staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CD101 antibody

Flow cytometric analysis of Human PBMC (human peripheral blood mononuclear cell) cells labelling CD101 with ab325180 at 1/50 dilution (1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Cells were co-stained with CD11c conjugated to Alexa Fluor®488.
Gated on viable cells.



Immunocytochemistry/ Immunofluorescence - Anti-CD101 antibody [EPR28974-15] (ab325180)


CD101 Immunocytochemistry/ Immunofluorescence staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CD101 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Human PBMC (human peripheral blood mononuclear cell) cells labelling CD101 with ab325180 at 1/50 (9.94 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/ab150081 1000 2ug/ml dilution (Green).

Confocal image showing cytoplasmic staining in a subset of human PBMC (shown in green). The counterstain was observed in magenta. Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

ab7291 Anti-alpha Tubulin mouse monoclonal antibody was used to counterstain tubulin at 1/ab7291 1000 1ug/ml dilution (Magenta). The Nuclear counterstain was DAPI (Blue).

-ve control 1: ab325180 at 1/50 dilution, followed by ab150120 Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) at 1/1000 dilution.
-ve control 2: ab7291 at 1/1000 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Flow Cytometry - Anti-CD101 antibody [EPR28974-15] (ab325180)

CD101 Flow Cytometry staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CD101 antibody

Flow cytometric analysis of Human PBMC (human peripheral blood mononuclear cell) cells labelling CD101 with ab325180 at 1/50 dilution (1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Cells were co-stained with CD8a conjugated to Pacific Blue.
Gated on viable cells.




Flow Cytometry - Anti-CD101 antibody [EPR28974-15] (ab325180)

CD101 Flow Cytometry staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CD101 antibody

Flow cytometric analysis of Human PBMC (human peripheral blood mononuclear cell) cells labelling CD101 with ab325180 at 1/50 dilution (1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Cells were co-stained with CD56 conjugated to PE.
Gated on viable cells.




Flow Cytometry - Anti-CD101 antibody [EPR28974-15] (ab325180)

CD101 Flow Cytometry staining of Human PBMC (human peripheral blood mononuclear cell) using rabbit Anti-CD101 antibody

Flow cytometric analysis of Human PBMC (human peripheral blood mononuclear cell) cells labelling CD101 with ab325180 at 1/50 dilution (1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Cells were co-stained with CD4 conjugated to Brilliant Violet 421.
Gated on viable cells.



Dot Blot - Anti-CD101 antibody [EPR28974-15] (ab325180)

CD101 Dot Blot staining using rabbit Anti-CD101 antibody

Dot blot analysis of CD101 using ab325180 at 1000 (0.497 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1:100,000 dilution.

Lane1: Human CD101 peptide
Lane2: Human IGSF3 peptide

Exposure time: 180 seconds.

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

This antibody does not cross-react with human IGSF3.

All lanes:
Dot Blot - Anti-CD101 antibody [EPR28974-15] (ab325180) at 1/1000 dilution

Lane 1:
Human CD101 peptide

Lane 2:
Human IGSF3 peptide

Secondary

All lanes:
Dot Blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.