abcam

BECKMAN COULTER

## ab326147

# SNv428 Anti-PKM antibody [EPR10138(B)]

Rabbit Recombinant Monoclonal PKM antibody - conjugated to SNv428. Suitable for Flow Cyt(Intra) and reacts with Human samples.

This is a Beckman Coulter product distributed by Abcam

Recombinant RabMAb

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Conjugation** | SNv428 |
| **Excitation/Emission** | Ex: 405nm, Em: 428nm |
| **Storage buffer** | pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA, 0.02% Pluronic F-68 |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | EPR10138(B) |
| **Purification technique** | Affinity purification Protein A |
| **Concentration** | 0.2 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/200 |
| **Notes** | - |

## Target data

See full target information PKM

**Function**

Catalyzes the final rate-limiting step of glycolysis by mediating the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP (PubMed:15996096, PubMed:1854723, PubMed:20847263). The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production (PubMed:15996096, PubMed:1854723, PubMed:20847263). The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and survival (PubMed:15996096, PubMed:1854723, PubMed:20847263). Isoform M2. Isoform specifically expressed during embryogenesis that has low pyruvate kinase activity by itself and requires allosteric activation by D-fructose 1,6-bisphosphate (FBP) for pyruvate kinase activity (PubMed:18337823, PubMed:20847263). In addition to its pyruvate kinase activity in the cytoplasm, also acts as a regulator of transcription in the nucleus by acting as a protein kinase (PubMed:18191611, PubMed:21620138, PubMed:22056988, PubMed:22306293, PubMed:22901803, PubMed:24120661). Translocates into the nucleus in response to various signals, such as EGF receptor activation, and homodimerizes, leading to its conversion into a protein threonine- and tyrosine-protein kinase (PubMed:22056988, PubMed:22306293, PubMed:22901803, PubMed:24120661, PubMed:26787900). Catalyzes phosphorylation of STAT3 at 'Tyr-705' and histone H3 at 'Thr-11' (H3T11ph), leading to activate transcription (PubMed:22306293, PubMed:22901803, PubMed:24120661). Its ability to activate transcription plays a role in cancer cells by promoting cell proliferation and promote tumorigenesis (PubMed:18337823, PubMed:22901803, PubMed:26787900). Promotes the expression of the immune checkpoint protein CD274 in BMAL1-deficient macrophages (By similarity). May also act as a translation regulator for a subset of mRNAs, independently of its pyruvate kinase activity: associates with subpools of endoplasmic reticulum-associated ribosomes, binds directly to the mRNAs translated at the endoplasmic reticulum and promotes translation of these endoplasmic reticulum-destined mRNAs (By similarity). Plays a role in caspase independent cell death of tumor cells (PubMed:17308100). Isoform M1. Pyruvate kinase isoform expressed in adult tissues, which replaces isoform M2 after birth (PubMed:18337823). In contrast to isoform M2, has high pyruvate kinase activity by itself and does not require allosteric activation by D-fructose 1,6-bisphosphate (FBP) for activity (PubMed:20847263).

## Storage

| | |
|---|---|
| **Shipped at conditions** | Blue Ice |
| **Appropriate short-term storage duration** | 1-2 weeks |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | +4°C |
| **Aliquoting information** | Upon delivery aliquot |
| **Storage information** | Avoid freeze / thaw cycle |

## Notes

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

**Patented technology**
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

This is a Beckman Coulter product distributed by Abcam.

The SuperNova dye included in this product is provided under an intellectual property license from Beckman Coulter Life Sciences. As this product contains the SuperNova dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity).

## Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

| | |
|---|---|
| **Activity summary** | PKM also known as pyruvate kinase muscle isozyme (PKM) and PEP is an enzyme that plays an important role in glycolysis by catalysing the conversion of phosphoenolpyruvate (PEP) to pyruvate yielding ATP in the process. The PKM protein has two isoforms PKM1 and PKM2 which result from alternative splicing of the PKM gene. The mass of PKM2 the more studied isoform is approximately 58 kDa. PKM is expressed in various tissues prominently in skeletal muscle heart brain and many tumor cells. Additionally PKM has significant activity in rapidly |

| | |
|---|---|
| | proliferating cells suggesting its importance in high-energy demanding environments. |
| **Biological function summary** | PKM functions not only in catalyzing the last step of glycolysis but also regulates metabolic and transcriptional processes. Specifically PKM2 is a participant in the regulation of gene expression and cellular response to oxidative stress and nutrient availability. It can exist as a dimer or tetramer with the latter being the more active form in glycolytic pathways while the dimeric form can translocate to the nucleus to perform functions unrelated to its glycolytic activity. These transformations make PKM part of a dynamic complex that responds to various cellular signals. |
| **Pathways** | PKM integrates into essential metabolic pathways including the glycolytic pathway and influences the pentose phosphate pathway. It works in conjunction with phosphofructokinase-1 (PFK1) another key glycolytic enzyme synchronizing the energy production process in cells. PKM2's non-metabolic roles involve interactions in signaling pathways related to cellular proliferation and survival often interacting with and modulating proteins like HIF-1α which plays a central role in cellular responses to hypoxia. |
| **Associated diseases and disorders** | PKM2 shows strong connections to cancer and metabolic diseases. Tumor cells often exhibit a shift in expression from PKM1 to PKM2 facilitating the altered metabolism known as the Warburg effect characterized by increased aerobic glycolysis. Its interaction with HIF-1α promotes adaptation to low oxygen environments typical in tumorous growth. Furthermore PKM disruptions or aberrations contribute to metabolic disorders such as diabetes where altered glucose metabolism becomes evident. The protein's behavior in these disease conditions indicates potential targets for therapeutic intervention highlighting the importance of PKM in both normal physiology and pathology. |

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

## 1 product image



### Flow Cytometry (Intracellular) - SNv428 Anti-PKM antibody [EPR10138(B)] (ab326147)

PKM Flow Cytometry (Intracellular) staining of Human peripheral mononuclear cells (PBMCs) using rabbit Anti-PKM antibody

Flow cytometry scatter plots showing human peripheral mononuclear cells (PBMCs), either stained with ab326147 (right) or FMO control (left). The cells were fixed and permeabilised using BD CytoFix/CytoPerm™ (20 min). The cells were incubated in 1x PBS containing 10 µg/ml human IgG, 10% normal goat serum and BD Pharmingen™ MonoBlock™ Leukocyte Staining Buffer (1/10) to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab326147) (1x 106 in 100µl at 1 µg/ml (1/200)) for 30min on ice. The cells were simultaneously stained with CD14. Acquisition of >30000 events were collected using a 100 mW Violet laser (405nm) and 450/45 bandpass filter.This data was acquired using a Beckman Coulter CytoFlexLX flow cytometer.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.