abcam

BECKMAN COULTER

ab326267

SNv428 Anti-GLS/Glutaminase antibody [EPR19525]

Rabbit Recombinant Monoclonal GLS antibody - conjugated to SNv428.
Suitable for Flow Cyt(Intra) and reacts with Human samples.

This is a Beckman Coulter product distributed by Abcam

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	SNv428
Excitation/Emission	Ex: 405nm, Em: 428nm
Storage buffer	pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA, 0.02% Pluronic F-68
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR19525
Purification technique	Affinity purification Protein A
Concentration	0.2 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200
Notes	-

Target data

See full target information GLS

Function	Catalyzes the first reaction in the primary pathway for the renal catabolism of glutamine. Plays a role in maintaining acid-base homeostasis. Regulates the levels of the neurotransmitter glutamate, the main excitatory neurotransmitter in the brain (PubMed:30239721, PubMed:30575854, PubMed:30970188). Isoform 2. Lacks catalytic activity.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply

- Animal-free batch production

For more information, read more on recombinant antibodies.

This is a Beckman Coulter product distributed by Abcam.

The SuperNova dye included in this product is provided under an intellectual property license from Beckman Coulter Life Sciences. As this product contains the SuperNova dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity).

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images



Flow Cytometry (Intracellular) - SNv428 Anti-GLS/Glutaminase antibody [EPR19525] (ab326267)


GLS/Glutaminase Flow Cytometry (Intracellular) staining using rabbit Anti-GLS/Glutaminase antibody

Flow cytometry scatter plots showing untreated human peripheral mononuclear cells (PBMCs) (top) and PBMCs activated with 5 µg/ml PHA for 18h (bottom), either stained with ab326267 (right) or FMO control (left). The cells were fixed and permeabilised using BD CytoFix/CytoPerm™ (20

min). The cells were incubated in 1x PBS containing 10 µg/ml human IgG, 10% normal goat serum and BD Pharmingen™ MonoBlock™ Leukocyte Staining Buffer (1/10) to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab326267) (1x 10^6 100µl at 1 µg/ml (1/200)) for 30min on ice. The cells were simultaneously stained with CD3.

Acquisition of >30000 events were collected using a 100 mW Violet laser (405nm) and 450/45 bandpass filter.

This data was acquired using a Beckman Coulter CytoFlexLX flow cytometer.



Flow Cytometry (Intracellular) - SNv428 Anti-GLS/Glutaminase antibody [EPR19525] (ab326267)

GLS/Glutaminase Flow Cytometry (Intracellular) staining using rabbit Anti-GLS/Glutaminase antibody

Flow cytometry overlay histogram showing left wild-type HAP1 positive cells and right negative GLS knockout HAP1 stained with ab326267 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab326267) (1x 10^6 in 100µl at 1 µg/ml (1/200)) for 30min at 22°C.

Unlabelled sample (black line) was also used as a control.

Acquisition of >5000 events were collected using a 100 mW Violet laser (405nm) and 450/45 bandpass filter.

This data was acquired using a Beckman Coulter CytoFlexLX flow cytometer.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.