abcam

ab326508

PE Anti-CD45RO antibody [UCHL1]

Key facts

Isotype	IgG2a
Host species	Mouse
Conjugation	PE
Excitation/Emission	Ex: 480;565nm, Em: 578nm
Storage buffer	pH: 7.4 Preservative: 0.098% Sodium azide Constituents: PBS, 0.2% BSA
Form	Liquid
Clonality	Monoclonal
Clone number	UCHL1
Purification technique	Size-exclusion chromatography
Specificity	The antibody UCHL1 recognizes an extracellular epitope of CD45R0, a 180 kDa low molecular weight isoform of the leukocyte common antigen (LCA). The antigen is expressed on a subset of memory/activated T cells and on cortical thymocytes.
Purification notes	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-

Notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Target data

See full target information PTPRC

Function	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor (PubMed:35767951). Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity (By similarity). Interacts with CLEC10A at antigen presenting cell-T cell contact; CLEC10A on immature dendritic cells recognizes Tn antigen-carrying PTPRC/CD45 receptor on effector T cells and modulates T cell activation threshold to limit autoreactivity. (Microbial infection) Acts as a receptor for human cytomegalovirus protein UL11 and mediates binding of UL11 to T-cells, leading to reduced induction of tyrosine phosphorylation of multiple signaling proteins upon T-cell receptor stimulation and impaired T-cell proliferation.

Storage

Shipped at conditions	Ambient - Can Ship with Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze, Store in the dark

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry - PE Anti-CD45RO antibody [UCHL1] (ab326508)

CD45RO Flow Cytometry staining of human peripheral whole blood using mouse Anti-CD45RO antibody


Flow cytometry surface staining pattern of human peripheral whole blood stained using ab326508 (20 µl reagent / 100 µl of peripheral whole blood).



Flow Cytometry - PE Anti-CD45RO antibody [UCHL1] (ab326508)

CD45RO Flow Cytometry staining of human lymphocytes using mouse Anti-CD45RO antibody

Flow cytometry multicolor surface staining of human lymphocytes stained using ab326508 (20 µl reagent / 100 µl of peripheral whole blood) and anti-human CD45RA (MEM-56) APC antibody (10 µl reagent / 100 µl of peripheral whole blood).



Flow Cytometry - PE Anti-CD45RO antibody [UCHL1] (ab326508)

CD45RO Flow Cytometry staining of human peripheral whole blood using mouse Anti-CD45RO antibody

Separation of human CD45R0 positive CD45RA negative lymphocytes (red-filled) from CD45R0 negative CD45RA positive lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using ab326508 (20 µl reagent / 100 µl of peripheral whole blood).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.