abcam

ab326879

Anti-CXCR6 antibody [EPR30957-585]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR30957-585
Purification technique	Affinity purification Protein A
Concentration	0.491 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

ICC/IF

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Mouse
Dilution info	-
Notes	-

WB

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information Cxcr6

Function	Receptor for the C-X-C chemokine CXCL16.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

5 product images




Flow Cytometry - Anti-CXCR6 antibody [EPR30957-585] (ab326879)

CXCR6 Flow Cytometry staining using rabbit Anti-CXCR6 antibody

Flow cytometric analysis of EL4 (mouse lymphoma T lymphocyte, Right) and Neuro-2a (mouse neuroblastoma neuroblast, Left) cells labelling CXCR6 with ab326879 at 1/500 dilution (0.1ug) / Red compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Negative control: Neuro-2a.




Flow Cytometry - Anti-CXCR6 antibody [EPR30957-585] (ab326879)

CXCR6 Flow Cytometry staining of Mouse spleen cell using rabbit Anti-CXCR6 antibody

Flow cytometric analysis of Mouse spleen cell cells labelling CXCR6 with ab326879 at 1/500 dilution (0.1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with NK1.1 conjugated to Brilliant Violet 421.




Flow Cytometry - Anti-CXCR6 antibody [EPR30957-585] (ab326879)

CXCR6 Flow Cytometry staining of Mouse spleen cell using rabbit Anti-CXCR6 antibody

Flow cytometric analysis of Mouse spleen cell cells labelling CXCR6 with ab326879 at 1/500 dilution (0.1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with B220 conjugated to Alexa Fluor®647.




Flow Cytometry - Anti-CXCR6 antibody [EPR30957-585] (ab326879)

CXCR6 Flow Cytometry staining of Mouse spleen cell using rabbit Anti-CXCR6 antibody

Flow cytometric analysis of Mouse spleen cell cells labelling CXCR6 with ab326879 at 1/500 dilution (0.1ug) / Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with CD3 conjugated to Alexa Fluor®647.



Flow Cytometry - Anti-CXCR6 antibody [EPR30957-585] (ab326879)

CXCR6 Flow Cytometry staining using rabbit Anti-CXCR6 antibody

Flow cytometric analysis of Mouse CD4+ PBMC cultured under Th1-polarizing conditions for 4 days (Upper left and right) and Mouse CD4+ PBMC cultured under Th2-polarizing conditions for 4 days (Lower left and right) cells labelling CXCR6 with ab326879 at 1/500 dilution (0.1ug) / Upper right and Lower right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells.
Cells were co-stained with CD25 conjugated to Brilliant Violet 421.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.