abcam

# ab326935

# Anti-TNF Receptor I antibody [EPR29704-145]

Recombinant | RabMAb | 20ul selling size

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | EPR29704-145 |
| **Purification technique** | Affinity purification Protein A |
| **Concentration** | 0.508 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

## Flow Cyt

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/500 |
| **Notes** | - |

## ICC/IF

**Tested**

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/50 |
| **Notes** | - |

## IHC-P

**Not recommended**

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |
| **Notes** | - |

## Target data

See full target information TNFRSF1A

| | |
|---|---|
| **Function** | Receptor for TNFSF2/TNF and homotrimeric TNFSF1/lymphotoxin-alpha. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Contributes to the induction of non-cytocidal TNF effects including anti-viral state and activation of the acid sphingomyelinase. |

## Storage

| | |
|---|---|
| **Shipped at conditions** | Conditional Ambient |
| **Appropriate short-term storage duration** | 1-2 weeks |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | -20°C |
| **Aliquoting information** | Upon delivery aliquot |
| **Storage information** | Avoid freeze / thaw cycle |

## Notes

**Patented technology**
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

## 3 product images




## Immunocytochemistry/ Immunofluorescence - Anti-TNF Receptor I antibody [EPR29704-145] (ab326935)


TNF Receptor I Immunocytochemistry/ Immunofluorescence staining of BT-549 (human Breast Mammary gland epithelial) using rabbit Anti-TNF Receptor I antibody

Immunofluorescent analysis of 100% methanol-fixed, 0.1% TritonX-100 permeabilized BT-549 (human Breast Mammary gland epithelial) and SH-SY5Y (human neuroblastoma epithelial cell) cells labelling TNF Receptor I with ab326935 at 1/50 (10.16 µg/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green).

Confocal image showing cytoplasmic staining in BT-549 cell line (shown in green). The counterstain was observed in magenta. Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Low expression: SH-SY5Y

ab7291 Anti-alpha Tubulin mouse monoclonal antibody was used to counterstain tubulin at 1/1000 dilution, followed by ab150120 Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) at 1/1000 dilution (Magenta).




## Flow Cytometry - Anti-TNF Receptor I antibody [EPR29704-145] (ab326935)

TNF Receptor I Flow Cytometry staining of BT-549 (human Breast, Mammary gland epithelial, Right) using rabbit Anti-TNF Receptor I antibody

Flow cytometric analysis of BT-549 (human Breast, Mammary gland epithelial, Right) and SH-SY5Y (human neuroblastoma epithelial cell, Left) cells labelling TNF Receptor I with ab326935 at 1/500 dilution (0.1ug) / Red compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells. Low expression: SH-SY5Y.



## Flow Cytometry - Anti-TNF Receptor I antibody [EPR29704-145] (ab326935)

TNF Receptor I Flow Cytometry staining of NCI-H1299 (human lung carcinoma epithelial cell, Right) using rabbit Anti-TNF Receptor I antibody

Flow cytometric analysis of NCI-H1299 (human lung carcinoma epithelial cell, Right) and 293T (human embryonic kidney epithelial cell, Left) cells labelling TNF Receptor I with ab326935 at 1/500 dilution (0.1ug) / Red compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Goat anti-Rabbit IgG (Alexa Fluor® 647, ab150083) at 1/5000 dilution was used as the secondary antibody.

Gated on viable cells. Low expression: 293T.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.