abcam

ab328536

PE Anti-CD207 antibody [2G3]

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	PE
Excitation/Emission	Ex: 480;565nm, Em: 578nm
Storage buffer	pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	Fusion protein of human CD207 extracellular part and IgG Fc fragment. Database link Q9UJ71
Clone number	2G3
Purification technique	Size-exclusion chromatography
Specificity	The mouse monoclonal antibody 2G3 recognizes an extracellular/luminal epitope of CD207, a transmembrane protein selectively expressed by Langerhans cells and by some other dendritic cell subsets. This antibody clone can target vaccine-peptides to dendritic cells.
Light chain type	kappa

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. Extracellular and intracellular staining.

Flow Cyt (Intra)

Expected

Species	Human
Dilution info	-
Notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. Extracellular and intracellular staining.

Target data

See full target information CD207

Function	Calcium-dependent lectin displaying mannose-binding specificity. Induces the formation of Birbeck granules (BGs); is a potent regulator of membrane superimposition and zippering. Binds to sulfated as well as mannosylated glycans, keratan sulfate (KS) and beta-glucans. Facilitates uptake of antigens and is involved in the routing and/or processing of antigen for presentation to T cells. Major receptor on primary Langerhans cells for Candida species, Saccharomyces species, and Malassezia furfur. Protects against human immunodeficiency virus-1 (HIV-1) infection. Binds to high-mannose structures present on the envelope glycoprotein which is followed by subsequent targeting of the virus to the Birbeck granules leading to its rapid degradation.

Storage

Shipped at conditions	Ambient - Can Ship with Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry - PE Anti-CD207 antibody [2G3] (ab328536)

CD207 Flow Cytometry staining of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells using mouse Anti-CD207 antibody


Flow cytometry multicolor surface staining pattern of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells stained using anti-human CD207 (2G3) PE antibody (10 µl reagent per milion cells in 100 µl of cell suspension) and anti-human CD11c (BU15) APC antibody (10 µl reagent per milion cells in 100 µl of cell suspension).



Flow Cytometry - PE Anti-CD207 antibody [2G3] (ab328536)

CD207 Flow Cytometry staining of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells using mouse Anti-CD207 antibody

Separation of human CD207 positive CD11c positive dendritic cells differentiated upon monocyte stimulation (GM-CSF + IL-4) (red-filled) from non-stimulated lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells stained using anti-human CD207 (2G3) PE antibody (10 µl reagent per milion cells in 100 µl of cell suspension).



Flow Cytometry - PE Anti-CD207 antibody [2G3] (ab328536)

CD207 Flow Cytometry staining of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells using mouse Anti-CD207 antibody

Flow cytometry surface staining pattern of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells stained using anti-human CD207 (2G3) PE antibody (10 µl reagent per milion cells in 100 µl of cell suspension).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.