abcam

ab328546

Alexa Fluor® 700 Anti-Ig Lambda Light Chain antibody [1-155-2]

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	Alexa Fluor® 700
Excitation/Emission	Ex: 702nm, Em: 723nm
Storage buffer	pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	Native Full Length Protein corresponding to Human Immunoglobulin lambda-1 light chain. Database link P0DOX8
Clone number	1-155-2
Purification technique	Size-exclusion chromatography
Specificity	The mouse monoclonal antibody 1-155-2 recognizes lambda light chains (22.5 kDa) of human immunoglobulin.
Light chain type	kappa

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. Extracellular and intracellular staining.

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µlof whole blood or 10^6 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. Extracellular and intracellular staining.

Target data

See full target information Immunoglobulin lambda-1 light chain

Function	Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:20176268, PubMed:22158414). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268).

Storage

Shipped at conditions	Ambient - Can Ship with Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Storage information	Do Not Freeze

Notes

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


3 product images



Flow Cytometry - Alexa Fluor® 700 Anti-Ig Lambda Light Chain antibody [1-155-2] (ab328546)

Ig Lambda Light Chain Flow Cytometry staining of human peripheral whole blood using mouse Anti-Ig Lambda Light Chain antibody


Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human Ig Lambda Light Chain (1-155-2) Alexa Fluor® 700 antibody (4 µl reagent / 100 µl of peripheral whole blood).



Flow Cytometry - Alexa Fluor® 700 Anti-Ig Lambda Light Chain antibody [1-155-2] (ab328546)

Ig Lambda Light Chain Flow Cytometry staining of human lymphocytes using mouse Anti-Ig Lambda Light Chain antibody

Flow cytometry multicolor surface staining of human lymphocytes stained using anti-human Ig Lambda Light Chain (1-155-2) Alexa Fluor® 700 antibody (4 µl reagent / 100 µl of peripheral whole blood) and anti-human CD19 (LT19) PE antibody (20 µl reagent / 100 µl of peripheral whole blood



Flow Cytometry - Alexa Fluor® 700 Anti-Ig Lambda Light Chain antibody [1-155-2] (ab328546)

Ig Lambda Light Chain Flow Cytometry staining of human peripheral whole blood using mouse Anti-Ig Lambda Light Chain antibody

Separation of human Ig Lambda Light Chain positive CD19 positive B cells (red-filled) from Ig Lambda Light Chain negative CD19 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human Ig Lambda Light Chain (1-155-2) Alexa Fluor® 700 antibody (4 µl reagent / 100 µl of peripheral whole blood).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.