abcam

**ab328558**

# PE Anti-CD187 antibody [10D1-J16]

## Key facts

| | |
|---|---|
| **Isotype** | IgG2a |
| **Host species** | Mouse |
| **Conjugation** | PE |
| **Excitation/Emission** | Ex: 480;565nm, Em: 578nm |
| **Storage buffer** | pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | Protein within Human ACKR3. Database link P25106 |
| **Clone number** | 10D1-J16 |
| **Purification technique** | Size-exclusion chromatography |
| **Specificity** | The mouse monoclonal antibody 10D1-J16 recognizes an extracellular epitope on CD187/CXCR7, a transmembrane protein of chemokine receptor family. |
| **Light chain type** | kappa |

## Reactivity data

## Flow Cyt

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |

| | |
|---|---|
| **Notes** | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |

## Target data

See full target information ACKR3

| | |
|---|---|
| **Function** | Atypical chemokine receptor that controls chemokine levels and localization via high-affinity chemokine binding that is uncoupled from classic ligand-driven signal transduction cascades, resulting instead in chemokine sequestration, degradation, or transcytosis. Also known as interceptor (internalizing receptor) or chemokine-scavenging receptor or chemokine decoy receptor. Acts as a receptor for chemokines CXCL11 and CXCL12/SDF1 (PubMed:16107333, PubMed:19255243, PubMed:19380869, PubMed:20161793, PubMed:22300987). Chemokine binding does not activate G-protein-mediated signal transduction but instead induces beta-arrestin recruitment, leading to ligand internalization and activation of MAPK signaling pathway (PubMed:16940167, PubMed:18653785, PubMed:20018651). Required for regulation of CXCR4 protein levels in migrating interneurons, thereby adapting their chemokine responsiveness (PubMed:16940167, PubMed:18653785). In glioma cells, transduces signals via MEK/ERK pathway, mediating resistance to apoptosis. Promotes cell growth and survival (PubMed:16940167, PubMed:20388803). Not involved in cell migration, adhesion or proliferation of normal hematopoietic progenitors but activated by CXCL11 in malignant hemapoietic cells, leading to phosphorylation of ERK1/2 (MAPK3/MAPK1) and enhanced cell adhesion and migration (PubMed:17804806, PubMed:18653785, PubMed:19641136, PubMed:20887389). Plays a regulatory role in CXCR4-mediated activation of cell surface integrins by CXCL12 (PubMed:18653785). Required for heart valve development (PubMed:17804806). Regulates axon guidance in the oculomotor system through the regulation of CXCL12 levels (PubMed:31211835). Acts as a receptor for SHLP2, mediating its effects on activation of proopiomelanocortin neurons in the arcuate nucleus of the hypothalamus which leads to suppression of food intake and increased energy expenditure (PubMed:37468558). (Microbial infection) Acts as a coreceptor with CXCR4 for a restricted number of HIV isolates. |

## Storage

| | |
|---|---|
| **Shipped at conditions** | Ambient - Can Ship with Ice |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | +4°C |
| **Storage information** | Do Not Freeze |

# Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**

We do not recommend this combination. It is not covered by our product promise.

---


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

## 1 product image




### Flow Cytometry - PE Anti-CD187 antibody [10D1-J16] (ab328558)

CD187 Flow Cytometry staining of MCF-7 cells using mouse Anti-CD187 antibody

Separation of MCF-7 cells stained using anti-human CD187 (10D1-J16) PE antibody (10 µl reagent per million cells in 100 µl of cell suspension, red-filled) from MCF-7 cells stained using mouse IgG2a isotype control (MOPC-173) PE antibody (concentration in sample 5 µg/ml, same as CD187 PE concentration, black-dashed) in flow cytometry analysis (surface staining) of MCF-7 cell suspension.

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.