abcam

ab51599

Anti-Cytokeratin 4 antibody [EP1599Y]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP1599Y
Purification technique	Affinity purification Protein A
Concentration	0.169 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

IP

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target data

See full target information KRT4

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


5 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599)

Cytokeratin 4 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 4 antibody


Immunohistochemical staining of paraffin embedded human cervical carcinoma tissue using ab51599 (1/100). Goat anti-rabbit HRP (1/2000) used for secondary staining.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599)

Cytokeratin 4 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Cytokeratin 4 antibody


ICC/IF image of ab51599 stained A431 cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab51599 at 1/100 dilution overnight at +4°C. The secondary antibody (pseudo-colored green) was Alexa Fluor® 488 goat anti- rabbit (ab150081) IgG (H+L) preadsorbed used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (pseudo-colored red) at a 1/200 dilution for 1h at room temperature. DAPI was used to stain the cell nuclei (pseudo-colored blue) at a concentration of 1.43µM for 1hour at room temperature.



Flow Cytometry (Intracellular) - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599)

Cytokeratin 4 Flow Cytometry (Intracellular) staining using rabbit Anti-Cytokeratin 4 antibody


Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Cytokeratin 4 with purified ab51599 at 1/100 dilution (10 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cells without incubation with primary antibody and secondary antibody (Blue).



Immunoprecipitation - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599)

Purified ab51599 at 1/50 dilution (2µg) immunoprecipitating Cytokeratin 4 in HaCaT whole cell lysate.
Lane 1 (input): HaCaT (Human skin keratinocyte) whole cell lysate 10µg
Lane 2 (+): ab51599 + HaCaT whole cell lysate.
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab51599 in HaCaT whole cell lysate.
VeriBlot for IP Detection Reagent (HRP) (ab131366) (1/1000 dilution) was used for Western blotting.
Blocking Buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM/TBST.
Observed band size: 59 kDa
All lanes:
Immunoprecipitation - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599)

Predicted band size: 57 kDa



Western blot - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599)

Cytokeratin 4 Western blot staining of A431 cell lysate using rabbit Anti-Cytokeratin 4 antibody

All lanes:
Western blot - Anti-Cytokeratin 4 antibody [EP1599Y] (ab51599) at 1/1000 dilution

All lanes:
A431 cell lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 57 kDa

Observed band size: 59 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.