abcam

# ab52600

# Anti-hnRNP K antibody [EP943Y]

Recombinant | RabMAb | 20ul selling size

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | EP943Y |
| **Purification technique** | Affinity purification Protein A |
| **Specificity** | This antibody detects both phosphorylated and unphosphorylated hnRNP K. |
| **Concentration** | 0.114 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

### IP

#### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/20 |
| **Notes** | **For unpurified use at 1/50** |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## WB

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100000 |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## ICC/IF

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/50 |
| **Notes** | **For unpurified use at 1/100 - 1/250.** |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/20 |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## IHC-P

### Tested

| | |
|---|---|
| **Species** | Mouse |
| **Dilution info** | 1/2000 |
| **Notes** | Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. |

| | |
|---|---|
| **Species** | Rat |
| **Dilution info** | 1/2000 |
| **Notes** | Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. |

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/2000 |
| **Notes** | Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. |

# Target data

See full target information HNRNPK

| | |
|---|---|
| **Function** | One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C) sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for pre-mRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA. Plays an important role in p53/TP53 response to DNA damage, acting at the level of both transcription activation and repression. When |

sumoylated, acts as a transcriptional coactivator of p53/TP53, playing a role in p21/CDKN1A and 14-3-3 sigma/SFN induction (By similarity). As far as transcription repression is concerned, acts by interacting with long intergenic RNA p21 (lincRNA-p21), a non-coding RNA induced by p53/TP53. This interaction is necessary for the induction of apoptosis, but not cell cycle arrest. As part of a ribonucleoprotein complex composed at least of ZNF827, HNRNPL and the circular RNA circZNF827 that nucleates the complex on chromatin, may negatively regulate the transcription of genes involved in neuronal differentiation (PubMed:33174841).

## Storage

| | |
|---|---|
| **Shipped at conditions** | Conditional Ambient |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | -20°C |
| **Storage information** | Stable for 12 months at -20°C |

## Notes

**Patented technology**
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

## Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

| | |
|---|---|
| **Activity summary** | Heterogeneous nuclear ribonucleoprotein K (hnRNP K) is a multifunctional protein that plays a critical role in RNA metabolism including RNA splicing translation and transcription. Alternate names for hnRNP K include synkaryon-associated protein and hnRNP KurB. It has a molecular mass of approximately 51 kDa and is ubiquitously expressed in the nucleus and cytoplasm of eukaryotic cells. This protein features three KH (K homology) domains that enable it to interact with nucleic acids and other proteins. |

| | |
|---|---|
| **Biological function summary** | HnRNP K serves as a transcriptional regulator and participates in the assembly of ribonucleoprotein complexes. As part of the ribonucleoprotein complex it interacts with both RNA and DNA influencing various gene expression pathways. It also modulates the activity of several transcription factors including c-Myc and has been implicated in cell proliferation and differentiation. This protein's versatile functions make it an important component in various cellular processes. |
| **Pathways** | HnRNP K plays an important role in signal transduction and the p53 pathway. In the p53 pathway hnRNP K acts as a coactivator enhancing the transcriptional activity of the tumor suppressor p53 and influencing cell cycle regulation and apoptosis. Additionally hnRNP K interacts with several proteins including the c-Myc oncogene to regulate gene expression associated with growth and cell cycle progression. |
| **Associated diseases and disorders** | HnRNP K has been linked to cancer and neurological disorders. Its interaction with c-Myc makes it relevant in various cancers where c-Myc is dysregulated contributing to unchecked cell proliferation. Furthermore alterations in hnRNP K expression or function have been associated with neurodegenerative disorders where it might interact with proteins like TDP-43 or FUS affecting neuronal function and survival. |

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**

We do not recommend this combination. It is not covered by our product promise.


---

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


## 11 product images



### Immunocytochemistry/ Immunofluorescence - Anti-hnRNP K antibody [EP943Y] (ab52600)

HnRNP K Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-hnRNP K antibody


Fluorescent immunostaining of HeLa cells using ab52600 at a dilution of 1/100-1/250.




### Immunocytochemistry/ Immunofluorescence - Anti-hnRNP K antibody [EP943Y] (ab52600)

HnRNP K Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-hnRNP K antibody


Immunocytochemistry/ Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling hnRNP K with purified ab52600 at 1:50 dilution (1.9 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with None. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-hnRNP K antibody [EP943Y] (ab52600)

HnRNP K Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-hnRNP K antibody


Immunohistochemical staining of paraffin-embedded human breast carcinoma tissue using ab52600 at a dilution of 1/100-1/250.




### Flow Cytometry (Intracellular) - Anti-hnRNP K antibody [EP943Y] (ab52600)

HnRNP K Flow Cytometry (Intracellular) staining using rabbit Anti-hnRNP K antibody


Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling hnRNP K with purified ab52600 at 1/20 dilution (10 µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).




## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-hnRNP K antibody [EP943Y] (ab52600)

HnRNP K Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-hnRNP K antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung carcinoma tissue sections labeling hnRNP K with purified ab52600 at 1:2000 dilution (0.05 µg/ml). Heat mediated antigen retrieval was performed Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




## Immunoprecipitation - Anti-hnRNP K antibody [EP943Y] (ab52600)

ab52600 at 1/100 immunoprecipitating hnRNP K in HeLa whole cell lysate.
For western blotting, a primary antibody was used at 1/1000 and a HRP-conjugated goat-anti-rabbit IgG was used as the secondary antibody (1/1000).


Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

All lanes:
Immunoprecipitation - Anti-hnRNP K antibody [EP943Y] (ab52600)

Predicted band size: 50 kDa


Observed band size: 60 kDa




## Immunoprecipitation - Anti-hnRNP K antibody [EP943Y] (ab52600)

ab52600 (purified) at 1:20 dilution (2µg) immunoprecipitating hnRNP K in Jurkat whole cell lysate.
Lane 1 (input): Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10µg
Lane 2 (+): ab52600 & Jurkat whole cell lysate
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab52600 in Jurkat whole cell lysate
For western blotting, VeriBlot for IP Detection Reagent (HRP) (ab131366) was used for detection at 1:1000 dilution.
Blocking and diluting buffer: 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-hnRNP K antibody [EP943Y] (ab52600)

Predicted band size: 50 kDa





## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-hnRNP K antibody [EP943Y] (ab52600)

HnRNP K Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-hnRNP K antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse stomach tissue sections labeling hnRNP K with purified ab52600 at 1:2000 dilution (0.05 µg/ml). Heat mediated antigen retrieval was performed Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was

used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-hnRNP K antibody [EP943Y] (ab52600)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat stomach tissue sections labeling hnRNP K with purified ab52600 at 1:2000 dilution (0.05 µg/ml). Heat mediated antigen retrieval was performed Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




## Western blot - Anti-hnRNP K antibody [EP943Y] (ab52600)

All lanes:
Western blot - Anti-hnRNP K antibody [EP943Y] (ab52600) at 1/100000 dilution


All lanes:
Jurkat cell lysate at 10 µg

Secondary

All lanes:
HRP-labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 50 kDa

Observed band size: ~60 kDa




## Western blot - Anti-hnRNP K antibody [EP943Y] (ab52600)

All lanes:
Western blot - Anti-hnRNP K antibody [EP943Y] (ab52600) at 1/10000 dilution

Lane 1:
Jurkat (Human T cell leukemia T lymphocyte) whole cell lysates at 20 µg

Lane 2:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates at 20 µg

Lane 3:
NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates at 20 µg

Lane 4:
MEF (Mouse embryonic fibroblast (immortalized)) whole cell lysates at 20 µg

Lane 5:
PC-12 (Rat adrenal gland pheochromocytoma ) whole cell lysates at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 50 kDa

Observed band size: 60 kDa

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.