abcam

ab52768

Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker

Lab Essentials | Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Tissue culture supernatant, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP1983Y
Purification technique	Affinity purification Protein A
Concentration	0.415 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	-

Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/5000 - 1/10000
Notes	-

Expected

Species	Rat, Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Rat
Dilution info	1/100 - 1/500
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Mouse
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/40 - 1/1000
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information MAP1LC3A

Function	Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes) (PubMed:20713600, PubMed:24290141). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation (PubMed:20713600). Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:31006537, PubMed:31006538).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


5 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker (ab52768)

MAP1LC3A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MAP1LC3A antibody

Immunohistochemical staining of MAP1LC3A in paraffin embedded human breast carcinoma using ab52768 at a 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker (ab52768)

MAP1LC3A Flow Cytometry (Intracellular) staining using rabbit Anti-MAP1LC3A antibody

Overlay histogram showing SHSY-5Y cells stained with ab52768 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab52768, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in SHSY-5Y cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.





Image from Buckingham EM et al. Biotechniques. 2014;57(5):241-4. Fig 1.; doi: 10.2144/000114226.

Immunocytochemistry/ Immunofluorescence - Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker (ab52768)


Immunocytochemistry/Immunofluorescence analysis of human keratinocytes (pannel A) and VZV-infected melanoma cells (panel E) labelling MAP1LC3A with ab52768.



Immunocytochemistry/ Immunofluorescence - Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker (ab52768)

MAP1LC3A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MAP1LC3A antibody

Immunocytochemistry/Immunofluorescence analysis of C6 (rat glioma) labelling MAP1LC3A with purified ab52768 at 1/500. Cells were fixed with 100% methanol. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody (ab150077). Nuclei counterstained with DAPI (blue).



Western blot - Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker (ab52768)

All lanes:
Western blot - Anti-MAP1LC3A antibody [EP1983Y] - Autophagosome Marker (ab52768) at 1/10000 dilution

All lanes:
Hela cell lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP labeled at 1/2000 dilution

Predicted band size: 14 kDa

Observed band size: 16 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.