abcam

ab52927

Anti-RON antibody [EP1132Y]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP1132Y
Purification technique	Affinity purification Protein A
Concentration	0.662 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Recombinant fragment, Recombinant full length protein - Human
Dilution info	-
Notes	-

IP

Expected

Species	Recombinant fragment
Dilution info	1/30
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

ELISA

Tested

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Recombinant fragment
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/10000
Notes	1/10000. Detects a band of approximately 35/150/180 kDa (predicted molecular weight: 185 kDa). See Notes section of Datasheet.

Not recommended

Species	Recombinant fragment, Recombinant full length protein - Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/20 - 1/40
Notes	Methanol fixed cells. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Not recommended

Species	Recombinant fragment, Recombinant full length protein - Human
Dilution info	-
Notes	-

Target data

See full target information MST1R

Function	Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to MST1 ligand. Regulates many physiological processes including cell survival, migration and differentiation. Ligand binding at the cell surface induces autophosphorylation of RON on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1 or the adapter GAB1. Recruitment of these downstream effectors by RON leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. RON signaling activates the wound healing response by promoting epithelial cell migration, proliferation as well as survival at the wound site. Also plays a role in the innate immune response by regulating the migration and phagocytic activity of macrophages. Alternatively, RON can also promote signals such as cell migration and proliferation in response to growth factors other than MST1 ligand.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

The major RON transcript is translated into a glycosylated single chain precursor, cleaved into a 185 kDa heterodimer (p185Ron) of 35 (alpha) and 150 kDa (beta) disulfide-linked chains, before exposure at the cell surface.
Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RON antibody [EP1132Y] (ab52927)

RON Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-RON antibody


Immunohistochemical staining of paraffin-embedded human stomach tissue using ab52927 at a dilution of 1/100-1/250.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-RON antibody [EP1132Y] (ab52927)


RON Flow Cytometry (Intracellular) staining using rabbit Anti-RON antibody

Overlay histogram showing THP1 cells stained with ab52927 (red line). The cells were fixed with methanol (5 min) and incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab52927 1/20 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (ab96899) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit monoclonal IgG (1µg/1x10^6 cells) used under the same conditions. Acquisition of >5000 events was performed. This antibody gave a decreased signal in THP1 cells fixed with 4% paraformaldehyde (10 min) used under the same conditions.Please note that Abcam do not have any data for use of this antibody in non-fixed cells. We welcome any customer feedback.



ELISA - Anti-RON antibody [EP1132Y] (ab52927)

ELISA analysis of Human RON recombinant protein at 1000 ng/mL with ab52927. An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) at 1/2500 dilution was used as the secondary antibody.



Western blot - Anti-RON antibody [EP1132Y] (ab52927)

RON Western blot staining of SKBR3 cell lysate using rabbit Anti-RON antibody

All lanes:
Western blot - Anti-RON antibody [EP1132Y] (ab52927) at 1/10000 dilution

All lanes:
SKBR3 cell lysate at 10 µg

Secondary

All lanes:
HRP-labelled goat anti-rabbit

Predicted band size: 152 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.