abcam

ab54190

PE Anti-Integrin beta 3 antibody [HM beta 3.1]

Anti-Integrin beta 3 antibody [HM beta 3.1] conjugated to PE - suitable for flow cytometry, and reacts with mouse, rat samples. Clone [HM beta 3.1] is a Armenian Hamster monoclonal antibody cited in 3 publications.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG
Host species	Armenian hamster
Conjugation	PE
Excitation/Emission	Ex: 480;565nm, Em: 578nm
Storage buffer	pH: 7.4 Preservative: 0.09% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	Native Full Length Protein corresponding to Mouse Integrin beta-3. Database link O54890
Clone number	HM beta 3.1
Purification technique	Affinity purification Protein G

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	-

Notes	Use 10µl of the suggested working dilution to label 10^6 cells in 100µl.

Expected

Species	Rat
Dilution info	1/10
Notes	-

Target data

See full target information Integrin beta-3

Function

Integrin alpha-V/beta-3 (ITGAV:ITGB3) is a receptor for cytotactin, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin alpha-IIB/beta-3 (ITGA2B:ITGB3) is a receptor for fibronectin, fibrinogen, plasminogen, prothrombin, thrombospondin and vitronectin. Integrins alpha-IIB/beta-3 and alpha-V/beta-3 recognize the sequence R-G-D in a wide array of ligands. Integrin alpha-IIB/beta-3 recognizes the sequence H-H-L-G-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin alpha-IIB/beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen. This step leads to rapid platelet aggregation which physically plugs ruptured endothelial surfaces. Fibrinogen binding enhances SELP expression in activated platelets (PubMed:19332769). ITGAV:ITGB3 binds to fractalkine (CX3CL1) and acts as its coreceptor in CX3CR1-dependent fractalkine signaling. ITGAV:ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling. ITGAV:ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling. ITGAV:ITGB3 binds to FGF2 and this binding is essential for FGF2 signaling (By similarity). ITGAV:ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling (By similarity). ITGAV:ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling (By similarity). ITGAV:ITGB3 binds to IL1B and this binding is essential for IL1B signaling (By similarity). ITGAV:ITGB3 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (By similarity). ITGAV:ITGB3 acts as a receptor for fibrillin-1 (FBN1) and mediates R-G-D-dependent cell adhesion to FBN1 (By similarity). ITGAV:ITGB3 binds to the Lilrb4a/Gp49b receptor and enhances the Lilrb4a-mediated inhibition of mast cell activation (PubMed:11323698). ITGAV:ITGB3 also suppresses marginal zone B cell antibody production through its interaction with Lilrb4a (PubMed:24935931). In brain, plays a role in synaptic transmission and plasticity (PubMed:18549786, PubMed:29038237). Involved in the regulation of the serotonin neurotransmission, is required to localize to specific compartments within the synapse the serotonin receptor SLC6A4 and for an appropriate reuptake of serotonin (PubMed:29038237). Controls excitatory synaptic strength by regulating GRIA2-containing AMPAR endocytosis, which affects AMPAR abundance and composition (PubMed:18549786). ITGAV:ITGB3 act as a receptor for CD40LG (By similarity). ITGAV:ITGB3 acts as a receptor for IBSP and promotes cell adhesion and migration to IBSP (By similarity). Integrin ITGA2B:ITGB3 is also the receptor of the

	erythrocyte-specific ICAM4 ligand involved in heterotypic cell-cell adhesion between erythrocytes and activated platelets (By similarity).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

Clone HM beta 3.1 partially inhibits the binding of CD61 to fibronectin.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Integrin beta 3 also known as ITGB3 beta 3 integrin or CD41/CD61 is a cell surface receptor that plays a role in cell adhesion and signal transduction. It is a member of the integrin family characterized by a heterodimer structure formed with an alpha subunit. This integrin has a molecular mass of approximately 88 kDa. Integrin beta 3 is expressed on platelets megakaryocytes and several other cell types such as endothelial cells and certain tumor cells.
Biological function summary	Integrins serve as the mechanical link between the extracellular matrix and the cytoskeleton. Beta 3 integrin forms a complex with different alpha subunits most notably alpha IIb to function as a receptor for fibrinogen fibronectin and vitronectin. This receptor complex is essential for platelet aggregation and normal hemostasis. Dysregulation can lead to altered cellular responses impacting processes like immune response and tumor progression.
Pathways	Integrin beta 3 participates in the signaling of outside-in and inside-out pathways that are fundamental in cellular dynamics. Of particular significance are the integrin-mediated signaling pathways and the TGF-beta signaling pathway which influence cellular proliferation and differentiation. Integrin beta 3 interacts with proteins such as talin and kindlin which are vital for activation and clustering of integrins during cell signaling processes.
Associated diseases and disorders	Altered expression or function of integrin beta 3 is associated with bleeding disorders like Glanzmann's thrombasthenia where platelet aggregation is

impaired. Additionally it plays a role in the progression of certain cancers due to its involvement in cell migration and invasion. The interaction of integrin beta 3 with proteins such as VEGF and MMPs is significant in the context of tumor angiogenesis and metastasis.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - PE Anti-Integrin beta 3 antibody [HM beta 3.1] (ab54190)

Integrin beta 3 Flow Cytometry staining of mouse peripheral blood platelets using armenian hamster Anti-Integrin beta 3 antibody

Staining of mouse peripheral blood platelets with RPE conjugated Hamster anti Mouse CD61 antibody, clone HM beta 3.1 used neat. Isotype control is represented by the blue histogram. Use 10µl of the suggested working dilution to 1x10^6 cells in 100µl.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.