abcam

ab59711

FITC Anti-TLR2 antibody [T2.5]

Anti-TLR2 antibody [T2.5] conjugated to FITC - suitable for flow cytometry, and reacts with mouse and human samples. Clone [T2.5] is a mouse monoclonal antibody cited in 27 publications.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	FITC
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	Synthetic Peptide within Mouse Tlr2. The exact immunogen used to generate this antibody is proprietary information. Database link Q9QUN7
Clone number	T2.5
Purification technique	Affinity purification Protein G
Concentration	0.1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	0.2 µm filtered.

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	10 µL for 10^6 Cells
Notes	ab91356 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Species	Mouse
Dilution info	10 µL for 10^6 Cells
Notes	ab91356 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Target data

See full target information TLR2

Function

Cooperates with LY96 to mediate the innate immune response to bacterial lipoproteins and other microbial cell wall components. Cooperates with TLR1 or TLR6 to mediate the innate immune response to bacterial lipoproteins or lipopeptides (PubMed:17889651, PubMed:21078852). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. May also activate immune cells and promote apoptosis in response to the lipid moiety of lipoproteins (PubMed:10426995, PubMed:10426996). Recognizes mycoplasmal macrophage-activating lipopeptide-2kD (MALP-2), soluble tuberculosis factor (STF), phenol-soluble modulin (PSM) and B.burgdorferi outer surface protein A lipoprotein (OspA-L) cooperatively with TLR6 (PubMed:11441107). Stimulation of monocytes in vitro with M.tuberculosis PstS1 induces p38 MAPK and ERK1/2 activation primarily via this receptor, but also partially via TLR4 (PubMed:16622205). MAPK activation in response to bacterial peptidoglycan also occurs via this receptor (PubMed:16622205). Acts as a receptor for M.tuberculosis lipoproteins LprA, LprG, LpqH and PstS1, some lipoproteins are dependent on other coreceptors (TLR1, CD14 and/or CD36); the lipoproteins act as agonists to modulate antigen presenting cell functions in response to the pathogen (PubMed:19362712). M.tuberculosis HSP70 (dnaK) but not HSP65 (groEL-2) acts via this protein to stimulate NF-kappa-B expression (PubMed:15809303). Recognizes M.tuberculosis major T-antigen EsxA (ESAT-6) which inhibits downstream MYD88-dependent signaling (shown in mouse) (By similarity). Forms activation clusters composed of several receptors depending on the ligand, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. Forms the cluster TLR2:TLR6:CD14:CD36 in response to diacylated lipopeptides and TLR2:TLR1:CD14 in response to triacylated lipopeptides (PubMed:16880211). Required for normal uptake of M.tuberculosis, a process that is inhibited by M.tuberculosis LppM (By similarity).

Storage

Shipped at conditions	Blue Ice

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Toll-like receptor 2 (TLR2) also called CD282 is a protein that detects molecules from bacteria and fungi. It has a mass of about 90 kDa. TLR2 is expressed on the surface of immune cells such as monocytes macrophages and neutrophils. This receptor recognizes pathogen-associated molecular patterns (PAMPs) and activates the immune response.
Biological function summary	TLR2 initiates signaling pathways that lead to the production of cytokines and chemokines helping to control infection. It forms dimers with other TLRs like TLR1 or TLR6 to expand its recognition range of microbial components. TLR2's activity is important for balancing pathogen recognition and avoiding overactive immune response. On macrophages TLR2 activation enhances their phagocytic ability.
Pathways	TLR2 plays roles in innate immune signaling pathways and inflammation pathways. It contributes significantly to the NF-kB and MAPK pathways which regulate inflammation and immune response. TLR2 often cooperates with proteins like MyD88 and TRAF6 in these pathways leading to downstream activation of transcription factors and production of pro-inflammatory molecules.
Associated diseases and disorders	TLR2 is associated with inflammatory and infectious diseases including sepsis and tuberculosis. During tuberculosis interactions between TLR2 and its ligands trigger immune responses that are important for bacterial control. Additionally TLR2’s overactivation links to sepsis by causing excessive inflammation. Researchers study TLR2 for potential therapeutic targeting in these diseases assessing how modulation can improve outcomes or reduce detrimental inflammation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

2 product images




Flow Cytometry - FITC Anti-TLR2 antibody [T2.5] (ab59711)

TLR2 Flow Cytometry staining using mouse Anti-TLR2 antibody

Overlay histogram showing THP1 cells stained with ab59711 (red line). The cells were fixed with 80% methanol (5 min) and incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab59711, 1µg/1x10^6 cells) for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 FITC (1µg/1x10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in THP1 cells fixed with 4% paraformaldehyde used under the same conditions.

Please note that Abcam do not have any data for use of this antibody on non-fixed cells. We welcome any customer feedback.



Flow Cytometry - FITC Anti-TLR2 antibody [T2.5] (ab59711)

TLR2 Flow Cytometry staining using mouse Anti-TLR2 antibody

Flow cytometric analysis of RAW 264.7 (mouse macrophage cell line transformed with Abelson murine leukemia virus) cells staining TLR2 using ab59711 at a 10 µg/mL (blue). Red and black represent the isotype control, cells only.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.