abcam

ab64085

Anti-TMEM16A antibody [SP31]

Anti-TMEM16A antibody [SP31] (ab64085) is a rabbit monoclonal antibody detecting TMEM16A in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human**.

- Biophysical QC for unrivalled batch-batch consistency
- Over 50 publications
- Trusted since 2008

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP31
Purification technique	Affinity purification Protein A
Concentration	0.055 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	PMID: 22685202.

Predicted

Species	Cynomolgus monkey
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	-

Predicted

Species	Cynomolgus monkey
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Cynomolgus monkey
Dilution info	-

Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Cynomolgus monkey
Dilution info	-
Notes	-

Target data

See full target information ANO1

Function	Calcium-activated chloride channel (CaCC) (PubMed:20056604, PubMed:22178883, PubMed:22946059, PubMed:32487539). Plays a role in transepithelial anion transport and smooth muscle contraction. Required for the normal functioning of the interstitial cells of Cajal (ICCs) which generate electrical pacemaker activity in gastrointestinal smooth muscles. Acts as a major contributor to basal and stimulated chloride conductance in airway epithelial cells and plays an important role in tracheal cartilage development. Required for CFTR activation by enhancing endoplasmic reticulum Ca(2+) store release and is also required for CFTR membrane expression (PubMed:28963502). Required for basal and ATP-dependent mucus secretion in airways and intestine, probably by controlling exocytosis of mucus-filled granules by providing Ca(2+) to an apical signaling compartment (By similarity). Contributes to airway mucus expression induced by interleukins IL3 and IL8 and by the asthma-associated protein CLCA1 and is required for expression of mucin MUC5AC (PubMed:33026825). However, was shown in another study not to be required for MUC5AC expression (PubMed:31732694). Plays a role in the propagation of Ca(2+) waves in Kolliker's organ in the cochlea and contributes to the refinement of auditory brainstem circuitries prior to hearing onset (By similarity). In vomeronasal sensory neurons, modulates spontaneous firing patterns in the absence of stimuli as well as the firing pattern of pheromone-evoked activity (By similarity). Responsible for calcium-activated chloride channel activity in type I taste cells of the vallate papillae (By similarity). Acts as a heat sensor in nociceptive neurons (By similarity). In dorsal root ganglion neurons, plays a role in mediating non-histaminergic Mas-related G-protein coupled receptor (MRGPR)-dependent itching, acting as a downstream effector of MRGPRs (By similarity). In the developing brain, required for the Ca(2+)-dependent process

extension of radial glial cells (By similarity). Isoform 4. Calcium-activated chloride channel (CaCC). Contributes to calcium-activated chloride secretion in human sweat gland epithelial cells. Shows increased basal chloride permeability and decreased Ca(2+)-induced chloride permeability. Isoform 5. Calcium-activated chloride channel (CaCC). Shows increased sensitivity to intracellular Ca(2+).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-TMEM16A antibody [SP31] (ab64085) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human samples.

What is the molecular weight of TMEM16A?
Anti-TMEM16A [SP31] (ab64085) specifically detects a band for TMEM16A (UniProt: Q5XXA6) at a molecular weight of 114kDa.

Collaboration
Anti-TMEM16A [SP31] (ab64085) is a clone from the portfolio of Spring Bioscience (Roche) SP clones which have been optimised for immunohistochemistry (IHC).

Trusted by the scientific community
Anti-TMEM16A [SP31] (ab64085) was first used in a scientific publication in 2008 and has been cited over 50 times in peer-reviewed journals.

Other related products
We have a range of other formats of antibody clone [SP31] also available for your convenience: ab64085, Carrier free - ab198412, Alexa Fluor® 647 - ab267509, PE - ab267510, APC - ab319337, Alexa Fluor® 488 - ab319587, Alexa Fluor® 594 - ab319830, Alexa Fluor® 555 - ab319971, Alexa Fluor® 750 - ab321687

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	TMEM16A also known as ANO1 is a transmembrane protein functioning as a calcium-activated chloride channel. It plays a significant role in the regulation of membrane potential and fluid secretion. TMEM16A has a molecular mass of approximately 114 kDa. This protein is expressed in various tissues including airway epithelial cells intestinal epithelium and smooth muscle cells. Its expression is also noticeable in parts of the nervous system indicating its widespread physiological importance.
Biological function summary	TMEM16A contributes to processes such as epithelial fluid transport and smooth muscle contraction. It is an integral component of a calcium-sensitive chloride conductance. This channel regulates ion transport across epithelial cell membranes influencing fluid secretion in airways and gastrointestinal tracts. In blood vessels TMEM16A helps control vascular tone by mediating smooth muscle cell contraction which is dependent on intracellular calcium levels.
Pathways	TMEM16A interacts with key cellular signaling cascades especially the calcium signaling pathway. This interaction has critical implications for chloride conductance in epithelial cells and smooth muscle function. TMEM16A is linked with proteins like calcium/calmodulin-dependent protein kinase II (CaMKII) which further modulates its activity. Additionally TMEM16A is implicated in pathways regulating mucin secretion which involves other proteins like cystic fibrosis transmembrane conductance regulator (CFTR).
Associated diseases and disorders	TMEM16A plays a role in respiratory diseases like asthma and cystic fibrosis where dysregulation in chloride channels leads to abnormal mucus secretion and airway obstruction. It also relates to hypertension due to its role in vascular tone modulation. Research shows TMEM16A interacts with proteins like CFTR in cystic fibrosis affecting chloride transport and fluid balance. In hypertension its interaction with calcium-sensing proteins further complicates vascular responsiveness and pressure regulation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

7 product images




Immunohistochemistry - Anti-TMEM16A antibody [SP31] (ab64085)

Immunohistochemical analysis of formalin fixed paraffin embedded human GIST labelling TMEM16A with ab64085 at a concentration of 1.14 µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with a OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was performed with DISCOVERY cell conditioning solution (CC1) 100°C, pH 8.5 for 32 mins.

ab64085 anti-TMEM16A [SP31] was incubated for 16 mins at 37°C. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.

Customers are encouraged to optimise antigen retrieval conditions, antibody concentration, incubation times and temperature for best results in their own IHC assay workflow (automated and manual).




This image is taken from Upregulation of TMEM16A Protein in Bronchial Epithelial Cells by Bacterial Pyocyanin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TMEM16A antibody [SP31] (ab64085)

TMEM16A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TMEM16A antibody

TMEM16A detected in paraffin-embedded sections of human submucosal glands from CF patients and control samples.
TMEM16A expression was modest in non-CF submucosal glands of non-CF samples (b) but markedly increased in tissues from CF patients (d) with a particularly strong signal (f) in histologically altered glands (e). Magnification: X630 in a X400 in b-f. Images 4Aa and 4Bace show hematoxylin and eosin staining.


This image was generated from the hybridoma version.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TMEM16A antibody [SP31] (ab64085)

TMEM16A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TMEM16A antibody


Immunohistochemical analysis of Human Gastrointestinal Stromal Tumor tissue labelling TMEM16A with ab64085.
This image was generated from the hybridoma version.



Flow Cytometry (Intracellular) - Anti-TMEM16A antibody [SP31] (ab64085)

TMEM16A Flow Cytometry (Intracellular) staining using rabbit Anti-TMEM16A antibody

Intracellular Flow Cytometry analysis of PC-3 (Human prostate adenocarcinoma epithelial cell) cells labeling TMEM16A with purified ab64085. (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluorr® 488, ab150077) secondary antibody. Isotype control - Rabbit monoclonal IgG (ab172730) / Black. Unlabeled control - Unlabelled cells / blue.
This image was generated from the hybridoma version.




This image is taken from Upregulation of TMEM16A Protein in Bronchial Epithelial Cells by Bacterial Pyocyanin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TMEM16A antibody [SP31] (ab64085)

TMEM16A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TMEM16A antibody

ab64085 showing expression of TMEM16A in human surface epithelium tissue. TMEM16A staining was mostly absent (b) and sometimes scanty (c) in the respiratory epithelium lining bronchi or bronchioles from CF patients (c arrows). A weak expression was also detectable in microvessels of peri-bronchial connective tissue (b arrows). Magnification: X400.
This image was generated from the hybridoma version.




Immunocytochemistry/ Immunofluorescence - Anti-TMEM16A antibody [SP31] (ab64085)

TMEM16A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-TMEM16A antibody

ab64085 staining TMEM16A in PC-3 (human prostate adenocarcinoma epithelial cell) cells. The cells were fixed with 4% formaldehyde, permeabilized in 100% methanol. The cells were then incubated with ab64085 at 1/20 dilution, followed by secondary antibody ab150077 AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). ab195889, Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used for counterstain at 1/200 dilution (Red). Nuclear DNA was labelled in blue with DAPI.
Confocal image showing membranous and cytoplasmic staining in PC-3 cell line.

Low expression control: LnCaP(PMID: 29899325)

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Western blot - Anti-TMEM16A antibody [SP31] (ab64085)

Blocking and diluting buffer and concentration: 5% NFDM/TBST Exposure time: Lane 1: 20 seconds Lane 2: 10 seconds The observed molecular weights are consistent with PMID: 22685202.
All lanes:
Western blot - Anti-TMEM16A antibody [SP31] (ab64085) at 1/1000 dilution

Lane 1:
PC-3 (Human prostate adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Capan-1 (Human pancreas adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 114 kDa

Observed band size: 130 kDa, 260 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.