abcam

ab76234

Anti-p130 antibody [EP2141Y]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP2141Y
Purification technique	Affinity purification Protein A
Concentration	0.701 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/250 - 1/300
Notes	-

Species	Rat

Dilution info	1/250 - 1/300
Notes	-

Species	Human
Dilution info	1/250 - 1/300
Notes	-

IP

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/1000
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information RBL2

Function Key regulator of entry into cell division. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases KMT5B and KMT5C, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Probably acts as a transcription repressor by recruiting chromatin-modifying enzymes to promoters. Potent inhibitor of E2F-mediated trans-activation, associates preferentially with E2F5. Binds to cyclins A and E. Binds to and may be involved in the transforming capacity of the adenovirus E1A protein. May act as a tumor suppressor.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Aliquoting information	Upon delivery aliquot
Storage information	Stable for 12 months at -20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	P130 also known as RBL2 (retinoblastoma-like 2) is a member of the retinoblastoma (Rb) protein family which includes pRb and p107. This protein has a molecular weight of about 130 kDa. It plays roles in regulating cell cycle by binding to E2F transcription factors consequently suppressing cell-cycle progression from G1 to S phase. p130 is ubiquitously expressed in various tissues but it shows higher levels in tissues with a low proliferation rate.
Biological function summary	The retinoblastoma-like 2 (p130) functions as an important regulator within the cell cycle checkpoint machinery. p130 forms a complex with E2F transcription factors selectively inhibiting their transcriptional activity. This inhibition plays an integral role in controlling cellular proliferation and ensuring that DNA replication does not occur unchecked. It also associates with cyclin-dependent kinase inhibitors further influencing the progression of the cell cycle.
Pathways	P130 is implicated in the control of the G1/S transition within the cell cycle pathway. Its function is closely linked with the Rb/E2F pathway which is critical for the regulation of cell proliferation. p130 collaborates with Rb and p107 within this pathway ensuring proper control of cell cycle transitions and maintenance of cellular growth.

Associated diseases and disorders	Retinoblastoma-like 2 (p130) shows significant connections to cancer and neurodegenerative disorders. Dysregulation of p130 along with its pathway partners like Rb can lead to unregulated cell growth contributing to oncogenesis. Additionally alterations in p130 function have implications in the development of Alzheimer's disease where cell cycle dysregulation occurs. Understanding p130's roles more deeply could provide insight into therapeutic targets for these conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


10 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human liver tissue sections labelling p130 with unpurified ab76234 at 1/250. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human liver tissue sections labelling p130 with purified ab76234 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labelling p130 with unpurified ab76234 at 1/50.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Flow Cytometry (Intracellular) staining using rabbit Anti-p130 antibody


Overlay histogram showing Jurkat cells stained with unpurified ab76234 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab76234 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluorr® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in Jurkat cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labelling p130 with unpurified ab76234 at 1/250. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat kidney tissue sections labelling p130 with unpurified ab76234 at 1/250. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labelling p130 with purified ab76234 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-p130 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat kidney tissue sections labelling p130 with purified ab76234 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Counterstained with hematoxylin.



Western blot - Anti-p130 antibody [EP2141Y] (ab76234)

Blocking buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM /TBST.

All lanes:
Western blot - Anti-p130 antibody [EP2141Y] (ab76234) at 1/1000 dilution


All lanes:
Daudi cell lysate at 10 µg

Secondary

All lanes:
Peroxidase-conjugated goat anti-rabbit IgG at 1/1000 dilution

Predicted band size: 128 kDa

Observed band size: 130 kDa



Western blot - Anti-p130 antibody [EP2141Y] (ab76234)

P130 Western blot staining of Daudi cell lysate using rabbit Anti-p130 antibody

Blocking buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM /TBST.

All lanes:
Western blot - Anti-p130 antibody [EP2141Y] (ab76234) at 1/1000 dilution

All lanes:
Daudi cell lysate at 10 µg

Secondary

All lanes:
Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 128 kDa

Observed band size: 130 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.