abcam

ab76295

Anti-HDAC3 antibody [EP462Y]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.05% Sodium azide Constituents: 99% Tissue culture supernatant, 0.1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP462Y
Purification technique	Affinity purification Protein A
Concentration	0.108 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Human
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

IP

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/2000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human

Dilution info	1/50
Notes	**For unpurified use at 1/100**

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/20
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information HDAC3

Function	Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates (PubMed:21030595, PubMed:21444723, PubMed:23911289, PubMed:25301942, PubMed:28167758, PubMed:28497810, PubMed:32404892, PubMed:22230954). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:23911289). Histone deacetylases act via the formation of large multiprotein complexes, such as N-Cor repressor complex, which activate the histone deacetylase activity (PubMed:23911289, PubMed:22230954). Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys-27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression (PubMed:23911289). Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (By similarity). Contributes, together with

XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:25190803). Regulates both the transcriptional activation and repression phases of the circadian clock in a deacetylase activity-independent manner (By similarity). During the activation phase, promotes the accumulation of ubiquitinated BMAL1 at the E-boxes and during the repression phase, blocks FBXL3-mediated CRY1/2 ubiquitination and promotes the interaction of CRY1 and BMAL1 (By similarity). The NCOR1-HDAC3 complex regulates the circadian expression of the core clock gene BMAL1 and the genes involved in lipid metabolism in the liver (By similarity). Also functions as a deacetylase for non-histone targets, such as KAT5, MEF2D, MAPK14, RARA and STAT3 (PubMed:15653507, PubMed:21030595, PubMed:21444723, PubMed:25301942, PubMed:28167758). Serves as a corepressor of RARA, mediating its deacetylation and repression, leading to inhibition of RARE DNA element binding (PubMed:28167758). In association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (PubMed:28167758). In addition to protein deacetylase activity, also acts as a protein-lysine deacylase by recognizing other acyl groups: catalyzes removal of (2E)-butenoyl (crotonyl), lactoyl (lactyl), 2-hydroxyisobutanoyl (2-hydroxyisobutyryl) and isonicotinyl acyl groups from lysine residues, leading to protein decrotonylation, delactylation, de-2-hydroxyisobutyrylation and deisonicotinylation, respectively (PubMed:28497810, PubMed:29192674, PubMed:34608293, PubMed:34545082, PubMed:35044827). Catalyzes decrotonylation of MAPRE1/EB1 (PubMed:34608293). Mediates delactylation NBN/NBS1, thereby inhibiting DNA double-strand breaks (DSBs) via homologous recombination (HR) (PubMed:38961290).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	HDAC3 or Histone Deacetylase 3 belongs to the class I histone deacetylases and plays an important mechanical role in epigenetic regulation by removing acetyl groups from histone proteins which causes chromatin condensation and gene expression modulation. Often referred to by its alternate name 3g6 HDAC3 has a molecular weight of approximately 49 kDa. The protein expresses broadly in various tissues including the brain liver and heart highlighting its widespread regulatory function across different body systems.
Biological function summary	HDAC3 acts as part of the nuclear receptor corepressor complex engaging with other subunits to modify chromatin structure and gene transcription. This protein influences various cellular processes such as cell cycle regulation and apoptosis. By altering chromatin architecture HDAC3 modulates the expression of numerous genes supporting cellular homeostasis and response to environmental signals.
Pathways	The HDAC3 protein integrates into key signaling networks including the Notch and NF-kB pathways. Within the Notch signaling pathway HDAC3 interacts with proteins like CSL modulating transcriptional responses critical for cell differentiation and developmental processes. In the NF-kB pathway it influences inflammation and immune responses by regulating the transcription of inflammatory genes showcasing its role in mediating complex signaling cascades.
Associated diseases and disorders	HDAC3 is pivotal in the development and progression of various diseases such as cancer and neurodegenerative disorders. The dysregulation of HDAC3 activity can lead to abnormal cell proliferation and impaired cell death contributing to tumor growth. In neurodegenerative conditions HDAC3 interacts with proteins like tau potentially influencing tau-related pathologies. HDAC3 inhibitors are under investigation for therapeutic approaches to these challenging medical conditions highlighting its relevance in disease management.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


8 product images



Immunocytochemistry/ Immunofluorescence - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-HDAC3 antibody


Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling HDAC3 with Purified ab76295 at 1:50 dilution (2.1 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488 ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC3 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung carcinoma tissue sections labeling HDAC3 with Purified ab76295 at 1:100 dilution (1.03 µg/ml). Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Flow Cytometry (Intracellular) - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Flow Cytometry (Intracellular) staining using rabbit Anti-HDAC3 antibody


Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling HDAC3 with Purified ab76295 at 1/20 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluorr® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC3 antibody


ab76295 at a 1/100-1/250 dilution staining human HDAC3 in breast carcinoma by Immunohistochemistry Formalin/PFA-fixed paraffin-embedded tissue.
Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC3 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat kidney tissue sections labeling HDAC3 with Purified ab76295 at 1:100 dilution (1.03 µg/ml). Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HDAC3 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse cerebrum tissue sections labeling HDAC3 with Purified ab76295 at 1:100 dilution (1.03 µg/ml). Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Western blot staining using rabbit Anti-HDAC3 antibody

All lanes:
Western blot - Anti-HDAC3 antibody [EP462Y] (ab76295) at 1/2000 dilution

Lane 1:
HeLa cell lysate at 20 µg

Lane 2:
Jurkat cell lysate at 20 µg

Lane 3:
HepG2 cell lysate at 20 µg

Lane 4:
NIH/3T3 cell lysate at 20 µg


Lane 5:
PC-12 cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 48 kDa

Observed band size: 49 kDa



Western blot - Anti-HDAC3 antibody [EP462Y] (ab76295)

HDAC3 Western blot staining of HeLa cell lysate using rabbit Anti-HDAC3 antibody

All lanes:
Western blot - Anti-HDAC3 antibody [EP462Y] (ab76295) at 1/2000 dilution

All lanes:
HeLa cell lysate at 10 µg

Secondary

All lanes:
goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 48 kDa

Observed band size: 49 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.