abcam

ab79393

Anti-MTCO2 antibody [EPR3314]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR3314
Purity	Tissue culture supernatant

Reactivity data

IP

Tested

Species	Human
Dilution info	1/50
Notes	-

WB

Tested

Species	Human
Dilution info	1/5000 - 1/10000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	5 µg/mL
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/60
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information MT-CO2

Function Component of the cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol-cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer

electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC) formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water molecules using 4 electrons from cytochrome c in the IMS and 4 protons from the mitochondrial matrix.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Stable for 12 months at -20°C

Notes

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	'MTCO2' also known as 'mt-co2' or 'mtco2e' is a mitochondrial gene that encodes for a component of the cytochrome c oxidase complex referred to as Complex IV in the electron transport chain. The protein plays a mechanical role in facilitating electron transfer within mitochondria an essential process in cellular respiration. MTCO2 is predominantly expressed in tissues with high energy demands such as muscle and neurons. The known mass of the MTCO2 protein is approximately 25 kDa. It sits in the mitochondrial inner membrane where it contributes to creating the proton gradient driving ATP synthesis.
Biological function summary	MTCO2 (or cytochrome c oxidase subunit II) serves as an important player in aerobic respiration. It is part of the cytochrome c oxidase complex which forms the last enzyme complex of the electron transport chain. As part of this complex MTCO2 facilitates the transfer of electrons from cytochrome c to oxygen resulting in the reduction of oxygen to water. This electron transfer is paired with proton translocation across the mitochondrial membrane which is critical for ATP production.
Pathways	MTCO2 contributes significantly to the oxidative phosphorylation pathway which is essential for ATP production in eukaryotic cells. It directly interacts with other components of the mitochondrial electron transport chain like cytochrome c and NADH dehydrogenase which are critical for maintaining the flow of electrons and the integrity of the energy production process. Another pathway it is part of is the apoptosis pathway regulated by non-lethal stress conditions where controlled release of cytochrome c can trigger programmed cell death.
Associated diseases and disorders	MTCO2 mutations and dysfunctions have been linked with mitochondrial disorders especially those affecting energy-demanding tissues leading to conditions such as mitochondrial myopathy and Leber's hereditary optic neuropathy. These disorders result from compromised oxidative phosphorylation leading to inadequate energy supply. The dysfunction of cytochrome c oxidase which contains the MTCO2 subunit is a central aspect of these diseases often tying this protein to other complexes within the electron transport chain that also underpin mitochondrial diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry (Intracellular) - Anti-MTCO2 antibody [EPR3314] (ab79393)

MTCO2 Flow Cytometry (Intracellular) staining using rabbit Anti-MTCO2 antibody


Overlay histogram showing HepG2 cells stained with ab79393 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab79393, 1/50 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (ab96899) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed.



Immunoprecipitation - Anti-MTCO2 antibody [EPR3314] (ab79393)

Purified ab79393 at 1/50 dilution (2µg) immunoprecipitating MTCO2 in HeLa whole cell lysate.
Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg
Lane 2 (+): ab79393 + HeLa whole cell lysate.
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab79393 in HeLa whole cell lysate.
VeriBlot for IP Detection Reagent (HRP) (ab131366) (1/1000 dilution) was used for Western blotting.
Blocking Buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM/TBST.
Observed band size: 21 kDa
All lanes:
Immunoprecipitation - Anti-MTCO2 antibody [EPR3314] (ab79393)

Predicted band size: 25 kDa



Western blot - Anti-MTCO2 antibody [EPR3314] (ab79393)

All lanes:
Western blot - Anti-MTCO2 antibody [EPR3314] (ab79393) at 1/10000 dilution

Lane 1:
K562 cell lysate at 10 µg

Lane 2:
MCF7 cell lysate at 10 µg

Lane 3:
THP1 cell lysate at 10 µg

Lane 4:
HeLa cell lysate at 10 µg

Secondary

All lanes:
goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 25 kDa

Observed band size: 21 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.