abcam

ab80681

Anti-IL-2 antibody [AP-MAB0844]

Key facts

Isotype	IgG2
Host species	Rat
Storage buffer	Constituents: PBS
Form	Lyophilized
Clonality	Monoclonal
Immunogen	Recombinant Full Length Protein corresponding to Mouse Il2. Database link P04351
Carrier free	Yes
Clone number	AP-MAB0844
Purification technique	Affinity purification Protein A
Reconstitution	reconstitute with PBS at 1mL
Purification notes	The IgG fraction of culture supernatant was purified by Protein A affinity chromatography.

Reactivity data

IP

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

FuncS (Neut/Block)

Expected

Species	Mouse
Dilution info	-
Notes	ab80681 can be used for neutralizing mouse IL2 in vitro.

IHC-Fr

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information Il2

Function

Cytokine produced by activated CD4-positive helper T-cells and to a lesser extend activated CD8-positive T-cells and natural killer (NK) cells that plays pivotal roles in the immune response and tolerance (PubMed:14614860, PubMed:9814585). Binds to a receptor complex composed of either the high-affinity trimeric IL-2R (IL2RA/CD25, IL2RB/CD122 and IL2RG/CD132) or the low-affinity dimeric IL-2R (IL2RB and IL2RG). Interaction with the receptor leads to oligomerization and conformation changes in the IL-2R subunits resulting in downstream signaling starting with phosphorylation of JAK1 and JAK3. In turn, JAK1 and JAK3 phosphorylate the receptor to form a docking site leading to the phosphorylation of several substrates including STAT5 (PubMed:14614860, PubMed:27018889). This process leads to activation of several pathways including STAT, phosphoinositide-3-kinase/PI3K and mitogen-activated protein kinase/MAPK pathways. Functions as a T-cell growth factor and can increase NK-cell cytolytic activity as well. Promotes strong proliferation of activated B-cells and subsequently immunoglobulin

production. Plays a pivotal role in regulating the adaptive immune system by controlling the survival and proliferation of regulatory T-cells, which are required for the maintenance of immune tolerance (PubMed:14614860). Moreover, participates in the differentiation and homeostasis of effector T-cell subsets, including Th1, Th2, Th17 as well as memory CD8-positive T-cells (PubMed:9814585).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Endotoxin Level: Extremely low level of LPS (< 0.002EU/µg IgG).
ab80681 was produced by *in vitro* culture, from a hybridoma (mouse myeloma fused with spleen cells from an immunized rat).

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.