abcam

ab90754

Anti-ENPP3/B10 antibody [NP4D6]

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	pH: 7.4 Preservative: 0.097% Sodium azide Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Cell preparation containing ENPP3 protein. The exact immunogen used to generate this antibody is proprietary information. Database link O14638
Clone number	NP4D6
Purification technique	Affinity purification Protein A
Specificity	The mouse monoclonal antibody NP4D6 reacts with CD203c, a transmembrane ectoenzyme expressed on basophils and mast cells, and overexpressed upon their activation.
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from cell culture supernatant with > 95% purity (by SDS-PAGE)

Reactivity data

Flow Cyt

Tested

Species	Human

Dilution info	1-4 µg/mL
Notes	-

Target data

See full target information ENPP3

Function	Hydrolase that metabolizes extracellular nucleotides, including ATP, GTP, UTP and CTP (PubMed:29717535, PubMed:9344668). Also hydrolyzes extracellular 2',3'-cGAMP (cyclic GMP-AMP), a second messenger that activates TMEM173/STING and triggers type-I interferon production, and is therefore involved in the regulation of innate immune response (PubMed:38749434). 2',3'-cGAMP appears to be further processed to GMP, AMP and inorganic phosphate (PubMed:38749434). Limits mast cells and basophils response during inflammation and during the chronic phases of allergic responses by eliminating extracellular ATP, a signaling molecule activating these cells in an autocrine manner. Metabolizes extracellular ATP in the lumen of the small intestine, and thereby prevents ATP-induced apoptosis of intestinal plasmacytoid dendritic cells (By similarity). Has a broad specificity and can also hydrolyze UDP-GlcNAc into UMP and GlcNAc-1-phosphate and potentially several other intracellular nucleotide sugars, including UDP-GalNAc, CMP-NeuAc, GDP-Fuc, and UDP-GlcA. Thereby, could modulate glycan biosynthesis and protein glycosylation (By similarity). Can hydrolyze extracellular dinucleoside polyphosphates, including the vasoactive adenosine polyphosphates as well (PubMed:12846830). In addition, displays an alkaline phosphodiesterase activity in vitro (PubMed:11342463).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The ENPP3/B10 protein also known as Ectonucleotide Pyrophosphatase/Phosphodiesterase 3 is an enzyme that plays a role in nucleotide

	metabolism by hydrolyzing phosphodiester and phosphosulfate bonds. This protein has a molecular mass of around 100 kDa. ENPP3 is mainly expressed on the surface of epithelial cells and you find it in tissues like the kidney liver and lung. In some contexts immune cells such as mast cells also show its expression.
Biological function summary	ENPP3 regulates the availability of extracellular nucleotides and phosphate metabolites. It is not typically considered part of a larger complex and rather contributes to various processes independently. By modulating concentrations of these extracellular molecules ENPP3 impacts cell signaling and nucleotide homeostasis which can affect cell proliferation and differentiation.
Pathways	This enzyme engages in the purinergic signaling pathway and can influence the MAPK/ERK pathway. Through its enzymatic action on nucleotides ENPP3 helps regulate signaling molecules which participate in processes such as cell growth and immune responses. Proteins related to these pathways include ATP-binding proteins like P2Y receptors and kinases involved in MAPK/ERK signaling.
Associated diseases and disorders	ENPP3's function links it to conditions such as allergy and asthma as its expression on mast cells affects histamine release. Changes in its activity are also noted in certain cancers where its role in signaling pathways can contribute to uncontrolled cell proliferation. In these disease contexts the protein interacts with other signaling molecules like cytokine receptors and oncogenic proteins which can modulate or exacerbate disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


2 product images



Flow Cytometry - Anti-ENPP3/B10 antibody [NP4D6] (ab90754)

ENPP3/B10 Flow Cytometry staining using mouse Anti-ENPP3/B10 antibody

Flow cytometry surface staining pattern of IgE stimulated human peripheral whole blood stained using ab90754 at 2 µg/ml.



Flow Cytometry - Anti-ENPP3/B10 antibody [NP4D6] (ab90754)

ENPP3/B10 Flow Cytometry staining using mouse Anti-ENPP3/B10 antibody

Separation of CD203c positive basophil granulocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of IgE stimulated human peripheral whole blood using ab90754 2 µg/ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.