abcam

ab90944

FITC Anti-CD14 antibody [47-3D6]

Anti-CD14 antibody [47-3D6] conjugated to FITC - suitable for flow cytometry, and reacts with human samples. Clone [47-3D6] is a mouse monoclonal antibody cited in 5 publications.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	FITC
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.2 Preservative: 0.1% Sodium azide Constituents: 0.9% BSA, 0.87% Sodium chloride, 0.328% Sodium phosphate
Form	Liquid
Clonality	Monoclonal
Immunogen	Native Full Length Protein corresponding to Human CD14. Database link P08571
Clone number	47-3D6
Purification technique	Affinity purification Immunogen
Specificity	ab90944 reacts with Human monocytes and macrophages; weak reactions may occur with neutrophils.
Light chain type	kappa

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	20 µL for 10^6 Cells
Notes	ab106163 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Target data

See full target information CD14

Function	Coreceptor for bacterial lipopolysaccharide (PubMed:1698311, PubMed:23264655). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:20133493, PubMed:22265692, PubMed:23264655). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:8612135). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed:16880211). Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed:23880187).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

Excitation wavelength: 488 nm Emission wavelength: 520 nm Molar composition: FITC/protein 6.0 – 8.0

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD14 also known as cluster of differentiation 14 is a glycoprotein with a molecular mass of approximately 55 kDa. This protein is expressed mainly on the surface of

	macrophages monocytes and to a lesser extent on neutrophils. CD14 exists in two forms: membrane-bound (mCD14) and soluble (sCD14). It functions as a co-receptor along with TLR4 (Toll-like receptor 4) and MD-2 for the detection of bacterial lipopolysaccharide (LPS) which is an important component of the outer membrane of Gram-negative bacteria.
Biological function summary	This glycoprotein plays a significant role in the innate immune system by recognizing pathogen-associated molecular patterns (PAMPs). It is part of a receptor complex where CD14 acts alongside LPS binding protein (LPS-R) TLR4 and MD-2 to facilitate pro-inflammatory signaling in response to microbial invasion. By binding to LPS CD14 initiates and amplifies the immune response leading to the production of cytokines and the recruitment of other immune cells to the site of infection.
Pathways	CD14 interacts with the toll-like receptor signaling pathway notably involving TLR4. Upon LPS detection CD14 assists in the activation of TLR4 triggering downstream signaling cascades such as NF-κB and MAPK pathways. These pathways result in the transcription of inflammatory cytokines and are essential for the host defense mechanism. The protein complex involving CD14 further interacts with adaptor proteins like MyD88 and TRIF facilitating signal propagation and the immune system's ability to respond to pathogens.
Associated diseases and disorders	Researchers have linked CD14 to sepsis and atherosclerosis. High levels of soluble CD14 (sCD14) are often observed in patients with sepsis indicating its participation in excessive systemic inflammation. In atherosclerosis CD14's role in recognizing LPS contributes to chronic inflammation and the development of plaques in arteries. Through these diseases CD14 connects with TLR4 emphasizing its impact on inflammatory responses and its potential as a therapeutic target.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - FITC Anti-CD14 antibody [47-3D6] (ab90944)

CD14 Flow Cytometry staining using mouse Anti-CD14 antibody

ab90944, at 20 µl per 10^6 cells, staining CD14 in Human normal peripheral blood by Flow cytometry.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.