abcam

ab92355

Anti-ILF3 antibody [EPR3626]

Anti-ILF3 antibody [EPR3626] (ab92355) is a rabbit monoclonal antibody detecting ILF3 in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency
- Over 20 publications
- Trusted since 2010

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR3626
Purification technique	Affinity purification Protein A
Concentration	0.6 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/10000
Notes	**For unpurified use at 1/100 - 1/250.** The use of an HRP/AP polymerized antibody is recommended as these have been shown to provide enhanced staining. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/10000 - 1/100000
Notes	-

Species	Rat
Dilution info	1/10000 - 1/100000
Notes	-

Species	Human
Dilution info	1/10000 - 1/100000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50 - 1/60

Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information ILF3

Function	RNA-binding protein that plays an essential role in the biogenesis of circular RNAs (circRNAs) which are produced by back-splicing circularization of pre-mRNAs. Within the nucleus, promotes circRNAs processing by stabilizing the regulatory elements residing in the flanking introns of the circularized exons. Plays thereby a role in the back-splicing of a subset of circRNAs (PubMed:28625552). As a consequence, participates in a wide range of transcriptional and post-transcriptional processes. Binds to poly-U elements and AU-rich elements (AREs) in the 3'-UTR of target mRNAs (PubMed:14731398). Upon viral infection, ILF3 accumulates in the cytoplasm and participates in the innate antiviral response (PubMed:21123651, PubMed:34110282). Mechanistically, ILF3 becomes phosphorylated and activated by the double-stranded RNA-activated protein kinase/PKR which releases ILF3 from cellular mature circRNAs. In turn, unbound ILF3 molecules are able to interact with and thus inhibit viral mRNAs (PubMed:21123651, PubMed:28625552). (Microbial infection) Plays a positive role

in HIV-1 virus production by binding to and thereby stabilizing HIV-1 RNA, together with ILF3.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Stable for 12 months at -20°C

Notes

What is this antibody validated in?
Anti-ILF3 antibody [EPR3626] (ab92355) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of ILF3?
Anti-ILF3 [EPR3626] (ab92355) specifically detects a band for ILF3 (UniProt: Q12906) at a molecular weight of 95kDa.

Trusted by the scientific community
Anti-ILF3 [EPR3626] (ab92355) was first used in a scientific publication in 2010 and has been cited over 20 times in peer-reviewed journals.

Other related products
We have a range of other formats of antibody clone [EPR3626] also available for your convenience: ab92355, HRP - ab206250, Carrier free - ab232546

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply

- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The ILF3 protein also known as Interleukin enhancer-binding factor 3 plays an important role in cellular mechanisms. It has a molecular mass of approximately 90 kDa and is expressed broadly in various tissues such as the liver kidney and spleen. ILF3 is part of the double-stranded RNA-binding proteins and it engages in regulating RNA metabolism processes. Scientists also refer to ILF3 as NF90 and it is known for its function in modulating gene expression by interacting with RNA.
Biological function summary	ILF3 interacts with RNA and proteins within the cell to control the stability and translation of mRNA. It acts as a component of various ribonucleoprotein complexes indicating its role in post-transcriptional regulation. ILF3 can bind to double-stranded RNA and single-stranded RNA motifs affecting the proliferation and differentiation of cells in which it is expressed. It also contributes significantly in cellular stress responses impacting how cells respond to environmental changes.
Pathways	ILF3 is involved in important cellular pathways such as RNA processing and antiviral defense mechanisms. In the RNA processing pathway ILF3 associates with proteins such as Drosha impacting microRNA biogenesis. Additionally in antiviral responses it identifies and binds viral RNAs initiating defensive cellular responses. These pathways position ILF3 as integral to maintaining cellular homeostasis during stress and infection.
Associated diseases and disorders	ILF3 has connections to cancer and viral infections. Aberrations in ILF3 expression or function are observed in certain cancers where it influences tumor progression by interacting with oncogenic pathways. In viral infections like those caused by hepatitis C virus ILF3 contributes to host defense by modulating immune responses. It interacts with proteins like RIG-I in the context of viral infections facilitating antiviral signaling pathways to control disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


6 product images



Flow Cytometry (Intracellular) - Anti-ILF3 antibody [EPR3626] (ab92355)

ILF3 Flow Cytometry (Intracellular) staining using rabbit Anti-ILF3 antibody


Overlay histogram showing HeLa cells stained with unpurified ab92355 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab92355, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (ab96899) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monclonal) (1µg/1x10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed.



Flow Cytometry (Intracellular) - Anti-ILF3 antibody [EPR3626] (ab92355)

ILF3 Flow Cytometry (Intracellular) staining using rabbit Anti-ILF3 antibody


Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling ILF3 with purified ab92355 at 1/60 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunocytochemistry/ Immunofluorescence - Anti-ILF3 antibody [EPR3626] (ab92355)

ILF3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ILF3 antibody


Immunofluorescence staining of HeLa cells with purified ab92355 at a working dilution of 1/500, counter-stained with DAPI. The secondary antibody was an Alexa Fluor® 488 conjugated goat anti-rabbit (ab150077), used at a dilution of 1/1000. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative control is shown in bottom right hand panel - for the negative control, PBS was used instead of the primary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ILF3 antibody [EPR3626] (ab92355)

ILF3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ILF3 antibody

ab92355 (unpurified) at 1/100 dilution staining ILF3 in paraffin-embedded Human kidney tissue, by immunohistochemistry.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-ILF3 antibody [EPR3626] (ab92355)

ILF3 Western blot staining using rabbit Anti-ILF3 antibody

Blocking and diluting buffer: 5% NFDM/TBST.
The two bands are reported by PMID: 22842455 and 27872311.
All lanes:
Western blot - Anti-ILF3 antibody [EPR3626] (ab92355) at 1/100000 dilution

Lane 1:
Raji (Human Burkitt's lymphoma B lymphocyte) whole cell lysates at 20 µg

Lane 2:
SH-SY5Y (Human neuroblastoma epithelial cell) whole cell lysates at 20 µg

Lane 3:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates at 20 µg


Lane 4:
Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysates at 20 µg

Lane 5:
C6 (Rat glial tumor glial cell) whole cell lysates at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 95 kDa



Western blot - Anti-ILF3 antibody [EPR3626] (ab92355)

All lanes:
Western blot - Anti-ILF3 antibody [EPR3626] (ab92355) at 1/100000 dilution

Lane 1:
Raji cell lysate at 10 µg

Lane 2:
HeLa cell lysate at 10 µg

Lane 3:
K562 cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit antibody at 1/2000 dilution

Predicted band size: 95 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.